

Please type a plus sign (+) inside this box

+

10/23/00

H
PTO/SB/05 (2/98)UTILITY
PATENT APPLICATION
TRANSMITTAL

Attorney Docket No.

PL08243.001 / 201130.407

First Inventor or Application Identifier

Nobuyuki Itoh

Title

HUMAN FGF-20 GENE AND GENE
EXPRESSION PRODUCTS

Express Mail Label No.

EL615483898US

Only for nonprovisional applications under 37 CFR § 1.53(b)

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

- 1.
- ☒
- General Authorization Form & Fee Transmittal
-
- (Submit an original and a duplicate for fee processing)

- 2.
- ☒
- Specification [Total Pages]
- 58**
-
- (preferred arrangement set forth below)
- Descriptive Title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention

- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

- 3.
- ☒
- Drawing(s) (35 USC 113) [Total Sheets]
- 17**

4. Declaration and Power of Attorney

2

[Total Pages]

- a.
- ☒
- Unsigned

- b.
- ☐
- Copy from a prior application (37 CFR 1.63(d))
-
- (for continuation/divisional with Box 17 completed)

- i.
- ☐
- DELETION OF INVENTOR(S)
-
- Signed statement attached deleting
-
- inventor(s) named in the prior
-
- application,
-
- see 37 CFR 1.63(d)(2) and 1.33(b)

- 5.
- ☐
- Incorporation By Reference (useable if box 4b is
-
- checked) The entire disclosure of the prior application,
-
- from which a copy of the oath or declaration is supplied
-
- under Box 4b, is considered to be part of the disclosure
-
- of the accompanying application and is hereby
-
- incorporated by reference therein.

- 6.
- ☐
- Microfiche Computer Program (Appendix)

7. Nucleotide and Amino Acid Sequence Submission
-
- (if applicable, all necessary)

- a.
- ☐
- Computer-Readable Copy
-
- b.
- ☐
- Paper Copy (identical to computer copy)
-
- c.
- ☐
- Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

- 8.
- ☐
- Assignment Papers (cover sheet & document(s))

- 9.
- ☐
- 37 CFR 3.73(b) Statement
-
- (when there is an assignee)
- ☒
- Power of Attorney

- 10.
- ☐
- English Translation Document (if applicable)

- 11.
- ☐
- Information Disclosure
-
- Statement (IDS)/PTO-1449
- ☐
- Copies of IDS
-
- Citations

- 12.
- ☐
- Preliminary Amendment

- 13.
- ☒
- Return Receipt Postcard

- 14.
- ☐
- Small Entity
-
- Statement(s)
- ☐
- Statement filed in prior application,
-
- Status still proper and desired

- 15.
- ☐
- Certified Copy of Priority Document(s)
-
- (if foreign priority is claimed)

- 16.
- ☒
- Other:
- Certificate of Express Mail
-
- Check No. 13629 for \$ 2,566.00

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information below and in a preliminary amendment



Continuation



Divisional



Continuation-In-Part (CIP)

of prior Application No.:

Prior application information: Examiner

Group / Art Unit

Claims the benefit of Provisional Application No. 60/161,162 filed October 22, 1999 and 60/187,856 filed March 8, 2000

CORRESPONDENCE ADDRESS

Jane E. R. Potter
Seed Intellectual Property Law Group PLLC
701 Fifth Avenue, Suite 6300
Seattle, Washington 98104-7092
Phone: (206) 622-4900 Fax: (206) 682-6031

Respectfully submitted,

TYPED or PRINTED NAME Jane E. R. Potter

SIGNATURE

REGISTRATION No. 33,332

Date

October 20, 2000

HUMAN FGF-20 GENE AND GENE EXPRESSION PRODUCTS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority from United States Patent Application

- 5 No. 60/161,162 filed October 22, 1999, and United States Patent Application No. 60,187,856 filed March 8, 2000, which are incorporated by reference herein in their entirety.

TECHNICAL FIELD

- 10 The present invention relates to nucleic acid sequences encoding members of the fibroblast growth factor (FGF) family, and to polypeptides encoded by the nucleic acid sequences.

BACKGROUND OF THE INVENTION

- 15 The substantia nigra is an area of the brain that has generated intensive research. Interest in the substantia nigra was originally based on the finding that degeneration of dopaminergic neurons in the area causes Parkinson's disease. In addition, the substantia nigra has been strongly implicated in thought and affective disorders (1). Therefore, neurotrophic factors for dopaminergic neurons in the substantia nigra are of substantial clinical interest.

- 20 Glial cell line-derived neurotrophic factor (GDNF) is the first neurotrophic factor documented to enhance the survival of midbrain dopaminergic neurons (Lin, L.-F.H. et al., *Science* 260:1130-1132 (1993)). Persephin, Artemin, BDGF and NT-3 also enhance the survival of midbrain dopaminergic neurons and have clinical potential in the treatment of Parkinson's disease (Milbrandt, J. et al., *Neuron* 20:245-253 (1998); Baloh, R.H. et al., *Neuron* 21:1291-1302 (1998); Hyman, C. et al., *Nature* 350:230-232 (1991); Hyman, C. et al., *J. Neurosci.* 14:335-347 (1994)). However, GDNF was reported to be widely expressed in neurons of the brain (Pochon, N.A. et al., *Eur. J. Neurosci.* 9:463-471 (1997)).
- 25 Persephin was also widely expressed in several major tissues including heart, kidney, liver

and brain (Milbrandt, J. et al., *Neuron* 20:245-253 (1998)). Artemin in brain was expressed in the basal ganglia and thalamus, suggesting that it influences the subcortical motor system (Baloh, R.H. et al., *Neuron* 21:1291-1302 (1998)). BDNF and NT-3 were predominantly expressed in the hippocampus (Ernfors, P. et al., *Neuron* 5:511-526 (1990)). Therefore, these neurotrophic factors appear not to be specific for dopaminergic neurons in the substantia nigra.

The prototypic fibroblast growth factors (FGFs), FGF-1 (aFGF) and FGF-2 (bFGF), were originally isolated from brain and pituitary as mitogens for fibroblasts. However, FGF-1 and FGF-2 are widely expressed in developing and adult tissues, and are polypeptides with multiple biological activities including angiogenesis, mitogenesis, cellular differentiation and repair of tissue injury (Baird, A. et al., *Cancer Cells* 3:239-243 (1991); Burgess, W.H. et al., *Annu. Rev. Biochem.* 58:575-606 (1989)). According to the published literature, the FGF family now consists of at least nineteen members, FGF-1 to FGF-19 (Dickson, C. et al., *Ann. NY Acad. Sci.* 638:18-26 (1991); Yoshida, T. et al., *Ann. NY Acad. Sci.* 638:27-37 (1991); Goldfarb, M. et al., *Ann. NY Acad. Sci.* 638:38-52 (1991); Coulier, F. et al., *Ann. NY Acad. Sci.* 638:53-61 (1991); Aaronson, S.A. et al., *Ann. NY Acad. Sci.* 638:62-77 (1991); Tanaka, A. et al., *Proc. Natl. Acad. Sci. USA* 89:8928-8932 (1992); Miyamoto, M. et al., *Mol. Cell. Biol.* 13:4251-4259 (1993); Yamasaki, M. et al., *J. Biol. Chem.* 271:15918-15921 (1996); Smallwood, P.M. et al., *Proc. Natl. Acad. Sci. USA* 93:9850-9857 (1996); McWhirter, J.R. et al., *Development* 124:3221-3232 (1997); Miyake, A. et al., *Biochem. Biophys. Res. Commun.* 243:148-152 (1998); Hoshikawa, M. et al., *Biochem. Biophys. Res. Commun.* 244:187-191 (1998); Ohbayashi, N. et al., *J. Biol. Chem.* 273:18161-18164 (1998); Nishimura, T. et al., *Biochim. Biophys. Acta* 1444:148-151 (1999)). FGF-3 was identified to be a common target for activation by the mouse mammary tumor virus (Dickson, C. et al., *Ann. NY Acad. Sci.* 638:18-26 (1991)). FGF-4 to FGF-6 were identified as oncogene products (Yoshida, T. et al., *Ann. NY Acad. Sci.* 638:27-37 (1991); Goldfarb, M. et al., *Ann. NY Acad. Sci.* 638:38-52 (1991); Coulier, F. et al., *Ann. NY Acad. Sci.* 638:53-61 (1991)). FGF-10 was identified from rat lung by homology-based polymerase chain reaction (PCR) (Yamasaki, M. et al., *J. Biol. Chem.* 271:15918-15921

(1996)). FGF-11 to FGF-14 (FGF homologous factors (FHF)s 1 to 4) were identified from human retina by a combination of random cDNA sequencing, data base searches and homology-based PCR (Smallwood, P.M. et al., *Proc. Natl. Acad. Sci. USA* 93:9850-9857 (1996)). FGF-15 was identified as a downstream target of a chimeric homeodomain oncoprotein (McWhirter, J.R. et al., *Development* 124:3221-3232 (1997)). FGF-16, FGF-17, and FGF-18 were identified from rat heart and embryos by homology-based PCR, respectively (Miyake, A. et al., *Biochem. Biophys. Res. Commun.* 243:148-152 (1998); Hoshikawa, M. et al., *Biochem. Biophys. Res. Commun.* 244:187-191 (1998); Ohbayashi, N. et al., *J. Biol. Chem.* 273:18161-18164 (1998)). Recently, FGF-19 was identified from human fetal brain by data base search (Nishimura, T. et al., *Biochim. Biophys. Acta* 1444:148-151 (1999)). They have a conserved ~120-amino acid residue core with ~30 to 60% amino acid identity. These FGFs also appear to play important roles in both developing and adult tissues. Thus, there is a need in the art for additional FGF molecules having functions and activities that differ from the known FGFs and for FGF molecules specifically expressed in regions of the brain implicated in human disease.

SUMMARY OF THE INVENTION

The present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising at least eight contiguous nucleotides of SEQ ID NO:1 or 3;
- (b) a polynucleotide that encodes a variant of the polypeptide encoded by (a); and
- (c) a polynucleotide encoding a protein expressed by a polynucleotide having the sequence of SEQ ID NO:1 or 3.

The invention further provides for the use of the isolated polynucleotides or fragments thereof as diagnostic probes or as primers.

The present invention also provides a composition comprising a polypeptide, wherein said polypeptide is selected from the group consisting of:

(a) a polypeptide comprising at least 6 contiguous amino acids encoded by SEQ ID NO:1 or 3;

(b) a polypeptide encoded by a polynucleotide comprising SEQ ID NO:1 or 3; and

5 (c) a variant of the polypeptide of (a) or (b).

Polypeptides of the invention are shown in SEQ ID NO:2 and 4. Other polypeptides comprise fragments of SEQ ID NO:2 and 4.

In certain preferred embodiments of the invention, the polynucleotide is operably linked to an expression control sequence. The invention further provides a host
10 cell, including bacterial, yeast, insect and mammalian cells, transformed with the polynucleotide sequence. The invention also provides full-length cDNA and full-length polynucleotides corresponding to SEQ ID NO:1 or 3.

Protein and polypeptide compositions of the invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody that
15 specifically reacts with such protein or polypeptide are also provided by the present invention.

The invention also provides for the production of large amounts of otherwise minor cell populations of cells to be used for generation of cDNA libraries for the isolation of rare molecules expressed in the precursors cell or progeny; cells produced
20 by treatment may directly express growth factors or other molecules, and conditioned media is screened in assays for novel activities.

The invention further provides for the isolation, self-renewal and survival of mammalian neural stem cells and the differentiation of their progeny.

The invention also provides for compositions and methods of preventing or
25 slowing degeneration of or increasing the numbers of dopaminergic neurons, such as in the substantia nigra, in disease states including Parkinson's disease.

The invention further provides for compositions and methods of preventing or slowing degeneration of, or for enhancing the growth of, cells in the inner ear.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Amino acid sequence comparison of rat FGF-20 with rat FGF-9 and FGF-16. Numbers refer to amino acid positions of FGF-9, FGF-16 and FGF-20. Asterisks indicate identical amino acid residues of the sequences.

5 Figure 2. The apparent evolutionary relationships of 20 members of the FGF family. The length of each horizontal line is proportional to the degree of amino acid sequence divergence. mFGF, rFGF, and hFGF indicate mouse FGF, rat FGF, and human FGF, respectively.

10 Figure 3. Localization of FGF-20 mRNA in rat brain. (A, B) Coronal sections were hybridized with ³⁵S-labeled FGF-20 antisense (A) and sense (B) probes, and exposed to X-ray film for 10 days. Section A is adjacent to section B. Scale bar = 0.5 cm. (C, D) Sections A and B were dipped in liquid emulsion and counterstained with Cresyl violet after 3 weeks. Dark-field photographs of the SNC in sections A and B are shown in C and D, respectively. White grains in the dark-field photograph show the localization of
15 FGF-20 mRNA. Scale bar = 50 μ m. SNC, substantial nigra pars compacta.

20 Figure 4. Expression of FGF-20 mRNA in rat brain. Rat brain poly (A)⁺RNA (10 μ g) was electrophoresed on a denaturing agarose gel (1%) containing formaldehyde and transferred onto a nitrocellulose membrane. Hybridization was performed with a ³²P labeled rat FGF-20 or β -actin cDNA probe. 28S and 18S indicate the positions of 28 and 18S rRNAs, respectively.

25 Figure 5. FGF-20 enhances survival of midbrain dopaminergic neurons. (A) Effect of FGF-20 on survival of midbrain dopaminergic neurons in serum-free medium. Midbrain cultured cells were incubated for 4 days in medium supplemented with 10% horse serum (control) or serum-free medium supplemented with FGF-20, and then the numbers of surviving dopaminergic neurons were determined. (B) Effect of FGF-20 for 24 h, and then treated with no (control) or 1mM glutamate for 10 min. The cultured cells were further incubated in the medium in the absence of glutamate and FGF-20 for 3 days, and then the numbers of surviving dopaminergic neurons were determined.

Figure 6. Figure 6 provides the DNA sequence (SEQ ID NO:1) of rat FGF-20.

Figure 7. Figure 7 provides the amino acid sequence (SEQ ID NO:2) of rat EGF-20.

Figure 8. Figure 8 provides the DNA (SEQ ID NO:3) and amino acid (SEQ ID NO:4) sequences of human FGF-20.

Figure 9. Figure 9 provides an alignment of the amino acid sequences of human (SEQ ID NO:4) and rat (SEQ ID NO:2) FGF-20.

Figure 10. Figure 10 provides codon usage for *E. coli*.

Figure 11. Figure 11 provides codon usage for yeast. The first field of information on each line of the table contains a three-letter code for an amino acid. The second field contains an unambiguous codon for that amino acid. The third field lists the number of occurrences of that codon in the genes from which the table is compiled. The fourth field lists the expected number of occurrences of that codon per 1,000 codons in genes whose codon usage is identical to that compiled in the codon frequency table. The last field contains the fraction of occurrences of the codon in its synonymous codon family.

Figure 12. Figure 12 provides codon usage for Drosophila.

DETAILED DESCRIPTION OF THE INVENTION

Because of their potent activities for promoting growth, proliferation, survival and differentiation of a wide variety of cells and tissue types, FGFs continue to be pursued as therapeutic agents for a number of different indications, including wound healing, such as musculo-skeletal conditions, for example, bone fractures, ligament and tissue repair, tendonitis, bursitis, etc.; skin conditions, for example, burns, cuts, lacerations, bed sores, slow healing ulcers, etc.; tissue protection and repair during myocardial infarction and ischemia, in the treatment of neurological conditions, for example, neurodegenerative disease and stroke, in the treatment of eye disease, including macular degeneration, and the like.

5
10

15

20

25

7

as shown in SEQ ID NO:1. A polynucleotide encoding the human FGF of the invention has the sequence as shown in SEQ ID NO:3. The rat polynucleotide was identified as encoding a member of the FGF family by the conserved regions throughout the amino acid sequence and by the regions of homology shared by the polynucleotides and genes encoding known FGF proteins.

The inventors believe that FGF-20 is a previously unidentified member of the FGF family. To date, over 19 human FGF proteins have been identified. In most cases, homologous proteins in the other mammals, particularly mice and rats, have also been identified. The human proteins vary to different degrees in terms of amino acid sequence, receptor specificity, tissue expression patterns, and biological activity.

The present FGF-20 differs in sequence from all the FGF proteins described to date in publications. FGF-20 shares some homology with FGF-9 and FGF-16.

As discussed herein, the knowledge about the roles played by various FGF proteins continues to grow, but is by far incomplete.

The present invention adds to this knowledge by disclosing that the FGF of SEQ ID NO:1 is highly expressed in dopaminergic neurons of the substantia nigra of brain, and human FGF-20 may play a role in development of and recovery from a neural disease, such as Parkinson's disease. FGF-20 is also preferentially expressed in rat embryo (E14.5) cochlea of the inner ear.

The invention therefore is based upon the identification, isolation, sequencing and expression patterns of a new fibroblast growth factor (FGF-20).

Isolation and Analysis of Rat cDNA encoding FGF-20 Members of the FGF family have a conserved ~ 120 –amino acid residue core with ~30 to 70% amino acid identity. Among the members of the FGF family, FGF-9 and FGF-16 are highly homologous (73% amino acid identity). According to the invention, DNA encoding a novel rat FGF has been identified. The nucleotide sequence of the entire coding region was determined by adaptor-ligation mediated polymerase chain reaction using rat-brain cDNA as a template and cassette-ligation mediated polymerase chain reaction using rat genomic DNA as a template. The nucleotide sequence of the coding region allowed for the

elucidation of the complete amino acid sequence of the FGF (212 amino acids), which has a conserved amino acid residue core (amino acids 62 to 197) (Fig. 1). Two cysteine residues that are well conserved in the FGF family are also conserved in the protein (amino acids 71 and 137) (Fig. 1). This protein is tentatively named FGF-20. FGF-20 is most similar to FGF-9 and FGF-16 (70 and 62% amino acid identity) among 19 members of the FGF family, respectively (Fig. 1). The apparent evolutionary relationships of twenty members of the FGF family are shown in Fig. 2. FGF-20 was closest to FGF-9 and FGF-16.

Expression of FGF-20 mRNA in Rat Tissues FGF-9 and FGF-16 mRNAs are preferentially expressed in rat kidney and heart, respectively (Miyamoto, M. et al., *Mol. Cell. Biol.* 13:4251-4259 (1993); Miyake, A. et al., *Biochem. Biophys. Res. Commun.* 243:148-152 (1998)). The expression of FGF-20 mRNA was examined in adult rat major tissues including brain, heart, lung, liver, kidney, and small intestine by polymerase chain reaction. FGF-20 mRNA was detected in the brain, but was undetectable or present in very low levels in other tissues. To confirm the expression of FGF-20 mRNA in rat brain, rat brain poly (A)⁺ RNA was examined by Northern blotting analysis using a ³²P-labeled rat FGF-20 cDNA probe. A faint but definite signal of FGF-20 mRNA was detected (Fig. 4). To confirm the integrity of the poly (A)⁺ RNA, the hybridized probe was washed from the membrane, and the membrane was rehybridized with a ³²P-labeled rat β -actin cDNA probe. A strong and discrete signal of β -actin mRNA was detected indicating that the poly (A)⁺ RNA was not degraded (Fig. 4). FGF-20 mRNA was also detected in rat embryos (E14.5), specifically in the cochlea of the inner ear, using ³⁵S-labeled FGF-20 anti-sense and sense cRNA probes (Fig. 10).

Expression of FGF-20 mRNA in Rat Brain To examine the expression of FGF-20 mRNA in rat brain, consecutive coronal sections of rat brain were analyzed by in situ hybridization with an ³⁵S-labeled antisense or sense FGF-20 cRNA probe. Discrete specific labeling was observed only in the substantia nigra pars compacta (Fig. 3A, C). No specific labeling was observed in other brain regions examined. The cellular localization of FGF-20 mRNA was examined by microscopy at higher magnification. By Nissle staining

of brain sections, glial cells can be identified as small intensely stained (dark) cells, while neurons are generally larger and less intensely stained (lighter) owing to their larger volume (Gerfen, C.R., *Methods in Neurosciences*, Academic Press, San Diego, CA, Vol. 1, pp. 79-97 (1989)). Black grains of labeled probes were found in most neurons of these brain areas (Fig. 3E). Dopaminergic neurons in the substantia nigra are preferentially localized in the substantia nigra pars compacta (Fallon, J.H. et al., *The Rat Nervous System*, 2nd Ed., Academic Press, San Diego, CA, pp. 215-238 (1995)). Furthermore, neurons in the substantia nigra pars compacta predominantly consist of dopaminergic neurons (Fallon, J.H. et al., *The Rat Nervous System*, 2nd Ed., Academic Press, San Diego, CA, pp. 215-238 (1995)). It is expected that FGF-20 is preferentially expressed in dopaminergic neurons in the substantia nigra pars compacta.

Preparation of Recombinant Rat FGF-20 To produce recombinant rat FGF-20, High Five insect cells were infected with recombinant baculovirus containing the rat FGF-20 cDNA with the 3'-terminal extension encoding E and His₆ tags. To detect recombinant FGF-20 in the culture medium, the medium was examined by Western blotting analysis with anti-E tag antibodies. A major band of 26.5 kDa was detected in the culture medium. The observed molecular mass of the major band was consistent with the calculated molecular mass of recombinant FGF-20 (26,247). This result indicates that FGF-20 is secreted, although on hydropathy plot analysis (Nielsen, H. et al., *Protein Engineering* 10:1-6 (1997)) the value of the amino-terminal region of FGF-20 was low, suggesting that FGF-20 has no signal sequence. Although FGF-9 and FGF-16 have no typical signal sequence in their amino termini, they are also secreted (Miyamoto, M. et al., *Mol. Cell. Biol.* 13:4251-4259 (1993), Miyake, A. et al., *Biochem. Biophys. Res. Commun.* 243:148-152 (1998)). Recombinant FGF-20 was purified from the culture medium by affinity chromatograph with Ni-NTA agarose and was analyzed by SDS-polyacrylamide gel electrophoresis under reducing conditions. A 26.5 kDa protein of FGF-20 was detected.

Neurotrophic Activity of FGF-20 for Rat Midbrain Dopaminergic Neurons FGFs are local signal molecules that act on proximal cells (Burgess, W.H. et al., *Annu. Rev. Biochem.* 58:575-606 (1989)). Therefore, it was expected that FGF-20 acts on

dopaminergic neurons in the substantia nigra in autocrine and/or paracrine manner. The neurotrophic activity of FGF-20 for cultured rat midbrain dopaminergic neurons was examined. When the dopaminergic neurons were cultured in a serum-free medium for 4 days, numbers of surviving dopaminergic neurons were greatly reduced. FGF-20 significantly enhanced survival of the dopaminergic neurons in the serum-free medium (Fig. 5A). The effect of FGF-20 on glutamate-induced neuronal death in cultured rat midbrain dopaminergic neurons was also examined. When the cultured cells were exposed to 1 mM glutamate for 10 min, the numbers of surviving dopaminergic neurons were reduced. FGF-20 also significantly enhanced survival of the dopaminergic neurons exposed to toxic concentrations of glutamate (Fig. 5B).

Several FGFs are expressed in brain and expected to play important roles as neurotrophic factors. FGF-1 and FGF-2 are abundant in brain (Gospodarowicz, D., *Methods Enzymol.* 147:106-119 (1987)) and exert survival enhancing effects on primary cultures from various regions of the brain (Walicke, P.A., *J. Neurosci.* 8:2618-2627 (1988)). FGF-1 is expressed predominantly in motor and sensory neurons of the midbrain and brainstem (Elde, R. et al., *Neuron* 7:349-364 (1991)). In contrast, FGF-2 is preferentially expressed in neurons in restricted regions including the cingulate cortex, indusium griemum, fasciola cinerum and hippocampus, and in astrocytes in widespread regions of the brain (Emoto, N. et al., *Growth Factors* 2:21-29 (1989); Woodward, W.R. et al., *J. Neurosci.* 12:142-152 (1992)). FGF-5 is weakly expressed in the cerebral cortex, hippocampus and thalamus (Haub, O. et al., *Proc. Natl. Acad. Sci. USA* 87:8022-8026 (1990)). FGF-9 and FGF-11 to FGF-14 are expressed in neurons of restricted regions including the hippocampus, thalamus, midbrain and brainstem (Yamamoto, S. et al., *Biochim. Biophys. Acta* 1398:38-41 (1998)). In contrast, FGF-20 of the invention was preferentially expressed in dopaminergic neurons of the substantia nigra. The expression profile of FGF-20 was quite distinct from those of other FGFs, indicating that FGF-20 plays a unique role in the brain.

Degeneration of dopaminergic neurons in the substantia nigra causes Parkinson's disease (Fallon, J.H. et al., *The Rat Nervous System*, 2nd Ed., Academic Press, San Diego, CA, pp. 215-238 (1995)). Therefore, neurotrophic factors for dopaminergic

neurons in the substantia nigra have received substantial attention. GDNF, Persephin, Artemin, BDNF, and NT-3 enhance survival of midbrain dopaminergic neurons (Lin, L.-F.H. et al., *Science* 260:1130-1132 (1993); Milbrandt, J. et al., *Neuron* 20:245-253 (1998); Baloh, R.H. et al., *Neuron* 21:1291-1302 (1998); Hyman, C. et al., *Nature* 350:230-232 (1991); Hyman, C. et al., *J. Neurosci.* 14:335-347 (1994)). However, their expression is not restricted to the substantia nigra (Pochon, N.A. et al., *Eur. J. Neurosci.* 9:463-471 (1997); Milbrandt, J. et al., *Neuron* 20:245-253 (1998); Baloh, R.H. et al., *Neuron* 21:1291-1302 (1998); Ernfors, P. et al., *Neuron* 5:511-526 (1990)). In contrast, the expression of FGF-20, which also enhanced the survival of midbrain dopaminergic neurons, was highly restricted in dopaminergic neurons in the substantia nigra pars compacta. Therefore, FGF-20 is expected to play an important role as a neurotrophic factor for dopaminergic neurons in the substantia nigra. It is therefore an important finding of the invention that FGF-20 is the first neurotrophic factor documented to be expressed preferentially in dopaminergic neurons of the substantia nigra.

It is believed that dopamine neurons are dysfunctional for, perhaps, years, before they are irreversibly damaged. (Dunnett, S.B. et al., *Nature* 399:A32-A39 (1999)) Thus, neurotrophic agents such as FGF-20 may be useful in preventing cell death or restoring function. The FGF-20 may be administered using gene transfer methods to block degeneration. Such methods have been used with neurotrophic factor GDNF (glial cell line-derived neurotrophic factor). In a rat Parkinson's model, nanogram amounts of BDNF and GDNF were measured from transduced cells, and the neuroprotective effect was in the order of 40-70% rescue of nigral dopamine neurons. Thus, transplants using fibroblasts or fibroblast cell lines engineered to secrete FGF-20 of the invention can allow secretion of the factor and rescue of nigral dopamine neurons. Alternatively, injection of the striatum or the substantia nigra region with viral vectors carrying the FGF-20 gene may also have a neuroprotective effect.

In Parkinson's Disease, neuronal degeneration in the substantia nigra generally is slow and protracted. This suggests that early intervention could block or slow down the degenerative process, perhaps up to 4 or 5 years before clinical symptoms appear.

A decline in striatal dopamine function can be detected by PET and SPECT imaging before the appearance of clinical symptoms, providing an opportunity for neuroprotective intervention at this early stage.

In vivo imaging of dopaminergic activity in the basal ganglia, using
 5 [18F]fluorodopa PET, can be used to monitor progress of the disease as well as the impact of treatment. A progressive reduction in fluorodopa signal is seen in brain tissue of pre-symptomatic and symptomatic individuals. After treatment with nigral tissue implant, the fluorodopa signal increases over time. (Dunnett et al., *supra*.) This and other techniques known in the art can be used to measure the effect of treatments described herein using
 10 FGF-20, and the clinician will be skilled in the art of determining appropriate treatment levels and regimens.

FGF-20 Expression in Rat Embryo Cochlea FGF-20 is preferentially expressed in the cochlea of the inner ear in rat embryos (E14.5). This supports a role for FGF-20 in the development and maintenance of normal ear function. Other previously-
 15 identified members of the FGF family contribute to normal ear growth and development. For example, sensory cells in the cochlea of the rat transiently express FGF-1 during the time of terminal innervation in the sensory epithelium (Dazert *et al.*, *J. Cell Physiol.* 177:123-129 (1998)). These authors also found that *in vitro*, spiral ganglion explants cultured in the presence of FGF-1 exhibited a dose-dependent increase in the number and
 20 length of neurites. In chick cochlea, FGF-1 mRNA levels increased in sensory epithelium of the cochlea in response to ototoxic damage, suggesting that the FGF system may be involved in the response of the cochlear epithelium to ototoxic damage. Pickles et al., *Dev. Neuroscience* 19:476-487 (1997). FGF-2 may help to regulate the proliferation step during hair cell development and regeneration after trauma in rats. Zheng et al., *J. Neuroscience*
 25 17:216-226 (1997). Thus, FGF molecules play several roles in maintaining normal development and function of the cochlea, and recovery of the cochlea from ototoxic damage. The absence of FGF receptor 3 contributes to inner ear defects in mice homozygous for skeletal and inner ear defects, including failure of pillar cell differentiation and tunnel of Corti formation, and profound deafness. Colvin et al., *Nat. Genet.* 12:390-

397 (1996). It is of interest that FGF-20 of the invention binds to FGF receptor 3c (Example 12). The fact that FGF-20 is expressed at a specific stage in rat inner ear development further suggests its importance in development of this tissue.

FGF-20 therefore may be suitable for treating a variety of conditions related to the ear. Currently, about 7.8 million Americans have mild hearing loss, 10 million have moderate hearing loss, and 2.7 million have profound or severe hearing loss. The causes include, but are not limited to, otosclerosis; Cogan's syndrome; Meniere's disease; Pendred's syndrome; diabetes-associated hearing loss (non-insulin-dependent diabetes mellitus in combination with obesity can cause tissue changes in the cochlea, McQueen et al., *J. Laryngol. Otol.* 113:113-118 (1999)); congenital malformations; autoimmune disease-related hearing loss; age-related hearing loss; deafness associated with lack of FGF receptor (Colvin et al., *Nat. Genet.* 12:390-397 (1996)); ischemia-related hearing disturbance; and other conditions in which cochlear structure and function plays a role. Administration of FGF-20 protein or polynucleotide may be used to treat inherited, congenital and acquired diseases of hearing and balance, by promoting the survival, proliferation or differentiation of cells of the inner ear.

Reference to FGF-20 herein is intended to be construed to include growth factors of any origin which are substantially homologous to and which are biologically equivalent to the FGF-20 characterized and described herein. Such substantially homologous growth factors may be native to any tissue or species and, similarly, biological activity can be characterized in any of a number of biological assay systems.

The term "biologically equivalent" is intended to mean that the compositions of the present invention are capable of demonstrating some or all of the same growth properties in a similar fashion, not necessarily to the same degree as the FGF-20 isolated as described herein or recombinantly produced human FGF-20 of the invention.

By "substantially homologous" it is meant that the degree of homology of human FGF-20 to FGF-20 from any species is greater than that between FGF-20 and any previously reported member of the FGF family.

Sequence identity or percent identity is intended to mean the percentage of same residues between two sequences, referenced to human FGF when determining percent identity with non-human FGF-20, referenced to FGF-20 when determining percent identity with non-FGF-20 growth factors, when the two sequences are aligned using the Clustal method (Higgins et al., *Cabios* 8:189-191 (1992)) of multiple sequence alignment in the Lasergene biocomputing software (DNASTAR, INC, Madison, WI). In this method, multiple alignments are carried out in a progressive manner, in which larger and larger alignment groups are assembled using similarity scores calculated from a series of pairwise alignments. Optimal sequence alignments are obtained by finding the maximum alignment score, which is the average of all scores between the separate residues in the alignment, determined from a residue weight table representing the probability of a given amino acid change occurring in two related proteins over a given evolutionary interval. Penalties for opening and lengthening gaps in the alignment contribute to the score. The default parameters used with this program are as follows: gap penalty for multiple alignment=10; gap length penalty for multiple alignment=10; k-tuple value in pairwise alignment=1; gap penalty in pairwise alignment=3; window value in pairwise alignment=5; diagonals saved in pairwise alignment=5. The residue weight table used for the alignment program is PAM250 (Dayhoff et al., in *Atlas of Protein Sequence and Structure*, Dayhoff, Ed., NDRF, Washington, Vol. 5, suppl. 3, p. 345, 1978).

Percent conservation is calculated from the above alignment by adding the percentage of identical residues to the percentage of positions at which the two residues represent a conservative substitution (defined as having a log odds value of greater than or equal to 0.3 in the PAM250 residue weight table). Conservation is referenced to human FGF-20 when determining percent conservation with non-human FGF-20, and referenced to FGF-20 when determining percent conservation with non-FGF-20 growth factors. Conservative amino acid changes satisfying this requirement are: R-K; E-D, Y-F, L-M; V-I, Q-H.

The invention provides FGF-20 proteins or variants thereof having one or more polymers covalently attached to one or more reactive amino acid side chains. By way

of example, not limitation, such polymers include polyethylene glycol (PEG), which can be attached to one or more free cysteine sulfhydryl residues, thereby blocking the formation of disulfide bonds and aggregation when the protein is exposed to oxidizing conditions. In addition, pegylation of FGF-20 proteins and/or muteins is expected to provide such improved properties as increased half-life, solubility, and protease resistance. FGF-20 proteins and/or muteins may alternatively be modified by the covalent addition of polymers to free amino groups such as the lysine epsilon or the N-terminal amino group. Preferred cysteines and lysines for covalent modification will be those not involved in receptor or heparin binding. In both human and rat FGF-20, the heparin binding site comprises amino acids 170-186. It will be apparent to one skilled in the art that the methods for assaying FGF-20 biochemical and/or biological activity may be employed in order to determine if modification of a particular amino acid residue affects the activity of the protein as desired.

It may be advantageous to improve the stability of FGF-20 by modifying one or more protease cleavage sites. Thus, the present invention provides FGF-20 variants in which one or more protease cleavage site has been altered by, for example, substitution of one or more amino acids at the cleavage site in order to create as FGF-20 variant with improved stability. Such improved protein stability may be beneficial during protein production and/or therapeutic use.

Suitable protease cleavage sites for modification are well known in the art and likely will vary depending on the particular application contemplated. For example, typical substitutions would include replacement of lysines or arginines with other amino acids such as alanine. The loss of activity, such as receptor binding or heparin binding, can be tested for as described herein.

FGF-20 can also include hybrid and modified forms of FGF-20 including fusion proteins and FGF-20 fragments and hybrid and modified forms in which certain amino acids have been deleted or replaced and modifications such as where one or more amino acids have been changed to a modified amino acid or unusual amino acid and modifications such as glycosylations so long as the hybrid or modified form retains the biological activity of FGF-20. By retaining the biological activity, it is meant that neuronal

survival is promoted, although not necessarily at the same level of potency as that of the FGF-20 isolated as described herein or that of the recombinantly produced human FGF-20. Fusion proteins can consist of the FGF-20 of the invention or fragment thereof and a signal sequence of a heterologous protein to promote secretion of the protein product.

5 Fusion proteins comprising FGF-20 or a biologically active or antigenic fragment thereof can be produced using methods known in the art. Such fusion proteins can be used therapeutically or can be produced in order to simplify the isolation and purification procedures. Histidine residues can be incorporated to allow immobilized metal affinity chromatography purification. Residues EQKLISEEDL contain the antigenic
10 determinant recognized by the myc monoclonal antibody and can be incorporated to allow myc monoclonal antibody-based affinity purification. A thrombin cleavage site can be incorporated to allow cleavage of the molecule at a chosen site; a preferred thrombin cleavage site consists of residues LVPRG. Purification of the molecule can be facilitated by incorporating a sequence, such as residues SAWRHPQFGG, which binds to
15 paramagnetic streptavidin beads. Such embodiments are described in WO 97/25345, which is incorporated by reference.

 The invention further includes chimeric molecules between FGF-20 and keratinocyte growth factor (KGF) (Reich-Slotky, R. et al., *J. Biol. Chem.* 270:29813-29818 (1995)). The chimeric molecule can contain specific regions or fragments of one or both of
20 the FGF-20 and KGF molecules, such as the FGF-20 fragments described below.

 The invention also includes fragments of FGF-20. Preferred fragments of SEQ ID NO:4 and 2 include: amino acids from about 170 to about 186; amino acids from about 1 to about 169; amino acids 2-211 (212 for SEQ ID NO:2); amino acids from about 1 to about 169 and about 187 to about 211 (212 for SEQ ID NO:2), wherein amino acids
25 about 169 and about 187 are joined by a peptide bond; and amino acids from about 59 to about 193. Such fragments can be prepared from the protein by standard biochemical methods or by expressing a polynucleotide encoding the fragment.

 FGF-20, or a fragment thereof, can be produced as a fusion protein comprising human serum albumin (HSA) or a portion thereof. Such fusion constructs are

suitable for enhancing expression of the FGF-20, or fragment thereof, in an eukaryotic host cell. Exemplary HSA portions include the N-terminal polypeptide (amino acids 1-369, 1-419, and intermediate lengths starting with amino acid 1), as disclosed in U.S. Patent No. 5,766,883, and publication WO 97/24445, incorporated by reference herein. Other
 5 chimeric polypeptides can include a HSA protein with FGF-20, or fragments thereof, attached to each of the C-terminal and N-terminal ends of the HSA. Such HSA constructs are disclosed in U.S. Patent No. 5,876,969, incorporated by reference herein.

Also included with the scope of the invention are FGF-20 molecules that differ from native FGF-20 by virtue of changes in biologically active sites. FGF-20 has a
 10 putative heparin binding site at amino acid residues 170-186. An FGF-20 molecule that does not bind heparin can be prepared by expressing DNA encoding FGF-20, wherein the corresponding codons for amino acid residues 170-186 have been deleted. Conversely, one or more additional heparin binding sites can be added to FGF-20 by, for example, expressing DNA encoding FGF-20 wherein the codons corresponding to residues 170-186
 15 are inserted at the desired position(s) in the reading frame. DNA encoding FGF-20 with altered receptor binding can likewise be produced. For example, it may be desirable to alter receptor specificity of FGF-20 by substituting the receptor binding regions of a different FGF for that of FGF-20.

Also included within the meaning of substantially homologous is any FGF-
 20 20 which may be isolated by virtue of cross-reactivity with antibodies to the FGF-20 described herein or whose encoding nucleotide sequences including genomic DNA, mRNA or cDNA may be isolated through hybridization with the complementary sequence of genomic or subgenomic nucleotide sequences or cDNA of the FGF-20 herein or fragments thereof. It will also be appreciated by one skilled in the art that degenerate DNA sequences
 25 can encode human FGF-20 and these are also intended to be included within the present invention, as are mammalian allelic variants of FGF-20.

Growth factors are thought to act at specific receptors. According to the invention, FGF-20 and as yet unknown members of this family of growth factors act through specific receptors having distinct distributions as has been shown for other growth

factor families. FGF-20 binds to FGF receptor 2 and FGF receptor 3, but does not bind to FGF receptor 1. Thus, its receptor binding profile differs from FGF-2 and FGF-4, which bind to FGF receptor 1.

A preferred hFGF-20 of the present invention has been identified and isolated in purified form as described. Also preferred is hFGF-20 prepared by recombinant DNA technology. By “pure form” or “purified form” or “substantially purified form” it is meant that an FGF-20 composition is substantially free of other proteins which are not FGF-20.

Recombinant human FGF-20 may be made by expressing the DNA sequences encoding FGF-20 in a suitable transformed host cell. Using methods well known in the art, the DNA encoding FGF-20 may be linked to an expression vector, transformed into a host cell and conditions established that are suitable for expression of FGF-20 by the transformed cell.

The DNA encoding FGF-20 can be engineered to take advantage of preferred codon usage of host cells. Codon usage in *Pseudomonas aeruginosa* is described in, for example, West et al., *Nucleic Acids Res.* 11:9323-9335 (1988). Codon usage in *Saccharomyces cerevisiae* is described in, for example, Lloyd et al., *Nucleic Acids Res.* 20:5289-5295 (1992). Codon preference in *Corynebacteria* and a comparison with *E. coli* preference is provided in Malubres et al., *Gene* 134:15-24 (1993). Codon usage in *Drosophila melanogaster* is described in, for example, Akashi, *Genetics* 136:927-935 (1994). Codon usage in yeast is also shown in Figure 11, and codon usage in *Drosophila* is shown in Figure 12.

Any suitable expression vector may be employed to produce recombinant human FGF-20 such as expression vectors for use in insect cells. Baculovirus expression systems can also be employed. A preferable method is expression in insect cells, such as Tr5 or Sf9 cells, using baculovirus vector.

The present invention includes nucleic acid sequences including sequences that encode human FGF-20. Also included within the scope of this invention are sequences that are substantially the same as the nucleic acid sequences encoding FGF-20. Such

Specific nucleic acid sequences can be modified by those skilled in the art and, thus, all nucleic acid sequences that code for the amino acid sequences of FGF-20 can likewise be so modified. The present invention thus also includes nucleic acid sequence which will hybridize with all such nucleic acid sequences or complements of the nucleic acid sequences where appropriate and encode a polypeptide having the neuronal cell survival promoting activities disclosed herein. The present invention also includes nucleic acid sequences that encode polypeptides that have neuronal cell survival promoting activity and that are recognized by antibodies that bind to FGF-20. Preferred methods and epitopes for raising antibodies are described in Example 10.

The present invention also encompasses vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the invention. This invention also includes host cells of any variety that have been transformed with vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the present invention.

Methods are also provided herein for producing FGF-20. Preparation can be by isolation from conditioned medium from a variety of cell types so long as the cell type produces FGF-20. A second and preferred method involves utilization of recombinant methods by isolating or obtaining a nucleic acid sequence encoding FGF-20, cloning the sequence along with appropriate regulatory sequences into suitable vectors and cell types, and expressing the sequence to produce FGF-20.

25 Although FGF-20 has been described on the basis of its ability to enhance
the survival of midbrain dopaminergic neurons, this factor may act on other cell types as
well. Thus, it is likely that FGF-20 can act on other neural cells.

It is also likely that FGF-20 will act on non-neuronal cells to promote their survival, growth or function. This expectation is based upon the activity of known growth

factors. Members of the FGF family act on many cell types of different function and embryologic origin.

The inventors herein have identified that FGF-20 is expressed in the brain, but not in other adult tissues, including heart, lung, liver, kidney and small intestine. This suggests a role for FGF-20 in, for example, Parkinson's disease and other diseases of neural tissue.

The present invention also includes therapeutic or pharmaceutical compositions comprising FGF-20 in an effective amount for treating patients with neuronal disease including Parkinson's disease, and a method comprising administering a therapeutically effective amount of FGF-20. These compositions and methods are useful for treating a number of diseases. The compositions and methods herein can also be useful to prevent degeneration and/or promote survival in other non-neuronal tissues as well. One skilled in the art can readily use a variety of assays known in the art to determine whether FGF-20 would be useful in promoting survival or functioning in a particular cell type, such as neuronal cells.

In certain circumstances, it may be desirable to modulate or decrease the amount of FGF-20 expressed. Thus, in another aspect of the present invention, FGF-20 anti-sense oligonucleotides can be made and a method utilized for diminishing the level of expression of FGF-20 by a cell comprising administering one or more FGF-20 anti-sense oligonucleotides. By FGF-20 anti-sense oligonucleotides reference is made to oligonucleotides that have a nucleotide sequence that interacts through base pairing with a specific complementary nucleic acid sequence involved in the expression of FGF-20 such that the expression of FGF-20 is reduced. Preferably, the specific nucleic acid sequence involved in the expression of FGF-20 is a genomic DNA molecule or mRNA molecule that encodes FGF-20. This genomic DNA molecule can comprise regulatory regions of the FGF-20 gene, or the coding sequence for mature FGF-20 protein. The term complementary to a nucleotide sequence in the context of FGF-20 antisense oligonucleotides and methods therefor means sufficiently complementary to such a sequence as to allow hybridization to that sequence in a cell, *i.e.*, under physiological conditions. The FGF-20 antisense

oligonucleotides preferably comprise a sequence containing from about 8 to about 100 nucleotides and more preferably the FGF-20 antisense oligonucleotides comprise from about 15 to about 30 nucleotides. The FGF-20 antisense oligonucleotides can also contain a variety of modifications that confer resistance to nucleolytic degradation such as, for example, modified internucleoside linkages (Uhlmann and Peyman, *Chemical Reviews* 90:543-548 1990; Schneider and Banner, *Tetrahedron Lett.* 31:335 (1990), which are incorporated by reference), modified nucleic acid bases and/or sugars and the like.

The therapeutic or pharmaceutical compositions of the present invention can be administered by any suitable route known in the art including for example intravenous, subcutaneous, intramuscular, transdermal, intrathecal or intracerebral. Administration can be either rapid as by injection or over a period of time as by slow infusion or administration of slow release formulation. For treating tissues in the central nervous system, administration can be by injection or infusion into the cerebrospinal fluid (CSF). When it is intended that FGF-20 be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of FGF-20 across the blood-brain barrier.

FGF-20 can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, FGF-20 can be coupled to any substance known in the art to promote penetration or transport across the blood-brain barrier such as an antibody to the transferrin receptor, and administered by intravenous injection (see, for example, Friden et al., *Science* 259:373-377 (1993), which is incorporated by reference). Furthermore, FGF-20 can be stably linked to a polymer such as polyethylene glycol to obtain desirable properties of solubility, stability, half-life and other pharmaceutically advantageous properties. (See, for example, Davis et al., *Enzyme Eng.* 4:169-73 (1978); Burnham, *Am. J. Hosp. Pharm.* 51:210-218 (1994), which are incorporated by reference.)

The compositions are usually employed in the form of pharmaceutical preparations. Such preparations are made in a manner well known in the pharmaceutical art. One preferred preparation utilizes a vehicle of physiological saline solution, but it is

contemplated that other pharmaceutically acceptable carriers such as physiological concentrations of other non-toxic salts, five percent aqueous glucose solution, sterile water or the like may also be used. It may also be desirable that a suitable buffer be present in the composition. Such solutions can, if desired, be lyophilized and stored in a sterile ampoule
 5 ready for reconstitution by the addition of sterile water for ready injection. The primary solvent can be aqueous or alternatively non-aqueous. FGF-20 can also be incorporated into a solid or semi-solid biologically compatible matrix which can be implanted into tissues requiring treatment.

The carrier can also contain other pharmaceutically-acceptable excipients for
 10 modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the carrier may contain still other pharmaceutically-acceptable excipients for modifying or maintaining release or absorption or penetration across the blood-brain barrier. Such excipients are those substances usually and customarily employed to formulate dosages for parenteral administration in either unit
 15 dosage or multi-dose form or for direct infusion into the cerebrospinal fluid by continuous or periodic infusion.

Dose administration can be repeated depending upon the pharmacokinetic parameters of the dosage formulation and the route of administration used.

It is also contemplated that certain formulations containing FGF-20 are to be
 20 administered orally. Such formulations are preferably encapsulated and formulated with suitable carriers in solid dosage forms. Some examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, gelatin, syrup, methyl cellulose, methyl- and
 25 propylhydroxybenzoates, talc, magnesium, stearate, water, mineral oil, and the like. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions may be formulated so as to provide rapid, sustained, or delayed release of the active ingredients after administration to the patient by employing procedures well known

in the art. The formulations can also contain substances that diminish proteolytic degradation and promote absorption such as, for example, surface active agents.

Depending on the treatment regimen contemplated, it may be desired to control the rate of release of FGF-20 protein or variant thereof to provide long-term treatment while minimizing the frequency of administration. Such treatment regimens may be desired, for example, where the FGF-20 protein is found to be relatively unstable such that the localized concentration of active protein is at an efficacious level for an insufficient period of time. Thus, for example, for certain diseases, it may not be desired or practical to perform repeated and frequent injections. The major advantages of such sustained release systems include targeted local delivery of drugs at a constant rate, less drug required to treat the disease state, minimization of possible side effects, and enhanced efficacy of treatment. Also, these forms of delivery systems are capable of protecting drugs that are unstable *in vivo* and that would normally require a frequent dosing interval. Under such circumstances, sustained release may be achieved by one of the methods readily available in the art such as the encapsulation of FGF-20 conjugated heparin-Sepharose beads to form heparin-alginate microspheres or the preparation of FGF-20 PLG microspheres.

Heparin-alginate microspheres have been successfully employed for the delivery of Basic Fibroblast Growth Factor to tissue (Lopez et al., *Journal of Pharmacology and Experimental Therapeutics* 282(1):385-390 (1997)). Similarly, Alginate/heparin-Sepharose microspheres and films have been used as drug carriers to control the release of a basic FGF-saponin conjugate in order to control its release in small doses. Addition of heparin to solutions of bFGF prevents losses in activity that accompany changes in pH or elevation in temperature. See, for example, Gospodarowicz et al., *J. Cell. Physiol.* 128:475-484 (1986).

As disclosed herein, FGF-20 has a heparin binding domain at residues 170-186. Accordingly, binding of FGF-20 to heparin may be employed in order to enhance its stability either during *in vivo* expression or administration or *in vitro* during various stages of protein purification. Thus, by the present invention, heparin may be added to a solution of FGF-20 and the activity assayed by the methods disclosed herein.

FGF-20 bound heparin-Sepharose beads may be encapsulated into calcium alginate microspheres to permit the controlled release of the heparin-stabilized FGF-20 protein. For example, microspheres may be constructed by dropping a mixed solution of sodium alginate with FGF-20 bound heparin-Sepharose beads into a hardening solution of calcium chloride. Spheres are formed instantaneously as the mixture enters the hardening solution. The size of the microsphere may be adjusted by passing the FGF-20 bound heparin-Sepharose beads through a cylinder of reduced cross-sectional area such as through a hypodermic needle.

Encapsulation efficiency may be determined by comparing the amount of encapsulated growth factor with that initially present in solution. For example, the FGF-20 may be stripped from the heparin-Sepharose beads with a solution of 3 M NaCl and functional activity assays may be performed.

The specific dose is calculated according to the approximate body weight or body surface area of the patient or the volume of body space to be occupied. The dose will also be calculated dependent upon the particular route of administration selected. Further refinement of the calculations necessary to determine the appropriate dosage for treatment is routinely made by those of ordinary skill in the art. Such calculations can be made without undue experimentation by one skilled in the art in light of the activity disclosed herein in assay preparations of target cells. Exact dosages are determined in conjunction with standard dose-response studies. It will be understood that the amount of the composition actually administered will be determined by a practitioner, in the light of the relevant circumstances including the condition or conditions to be treated, the choice of composition to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration.

In one embodiment of this invention, FGF-20 may be therapeutically administered by implanting into patients vectors or cells capable of producing a biologically-active form of FGF-20 or a precursor of FGF-20, *i.e.*, a molecule that can be readily converted to a biological-active form of FGF-20 by the body. In one approach cells that secrete FGF-20 may be encapsulated into semipermeable membranes for implantation

into a patient. The cells can be cells that normally express FGF-20 or a precursor thereof or the cells can be transformed to express FGF-20 or a precursor thereof. It is preferred that the cell be of human origin and that the FGF-20 be human FGF-20 when the patient is human. However, the formulations and methods herein can be used for veterinary as well
 5 as human applications and the term "patient" as used herein is intended to include human and veterinary patients.

Cells can be grown *ex vivo* for use in transplantation or engraftment into patients (Muench et al., *Leuk. & Lymph. 16*:1-11 (1994), which is incorporated by reference). In another embodiment of the present invention, FGF-20 is used to promote the
 10 *ex vivo* expansion of a cells for transplantation or engraftment. Current methods have used bioreactor culture systems containing factors such as erythropoietin, colony stimulating factors, stem cell factor, and interleukins to expand hematopoietic progenitor cells for erythrocytes, monocytes, neutrophils, and lymphocytes (Verfaillie, *Stem Cells 12*:466-476 (1994), which is incorporated by reference). These stem cells can be isolated from the
 15 marrow of human donors, from human peripheral blood, or from umbilical cord blood cells. The expanded blood cells are used to treat patients who lack these cells as a result of specific disease conditions or as a result of high dose chemotherapy for treatment of malignancy (George, *Stem Cells 12*(Suppl 1):249-255 (1994), which is incorporated by reference). In the case of cell transplant after chemotherapy, autologous transplants can be
 20 performed by removing bone marrow cells before chemotherapy, expanding the cells *ex vivo* using methods that also function to purge malignant cells, and transplanting the expanded cells back into the patient following chemotherapy (for review, see Rummel and Van Zant, *J. Hematotherapy 3*:213-218 (1994), which is incorporated by reference). Since FGF-20 is expressed in neural cells, it is believed that FGF-20 can function to prevent or
 25 slow the degeneration of dopaminergic neurons, such as substantia nigra.

In a number of circumstances it would be desirable to determine the levels of FGF-20 in a patient. The identification of FGF-20 along with the data herein showing expression of FGF-20 provides the basis for the conclusion that the presence of FGF-20 serves a normal physiological function related to cell growth and survival. Indeed, other

neurotrophic factors are known to play a role in the function of neuronal and non-neuronal tissues. (Scully and Otten, *Cell Bol. Int.* 19:459-469 (1995); Otten and Gadiant, *Int. J. Devl. Neurosciences* 13:147-151 (1995), which are incorporated by reference.) Endogenously produced FGF-20 may also play a role in certain disease conditions, particularly where there is cellular degeneration such as in neurodegenerative conditions or diseases. Other neurotrophic factors are known to change during disease conditions. For example, in multiple sclerosis, levels of NGF protein in the cerebrospinal fluid are increased during acute phases of the disease (Bracci-Laudiero et al., *Neuroscience Lett.* 147:9-12 (1992), which is incorporated by reference) and in systemic lupus erythematosus there is a correlation between inflammatory episodes and NGF levels in sera (Bracci-Laudiero et al., *NeuroReport* 4:563-565 (1993), which is incorporated by reference).

Given that FGF-20 is expressed in adult neural cells and in cochlear cells during embryonic development, it is likely that the level of FGF-20 may be altered in a variety of conditions and that quantification of FGF-20 levels would provide clinically useful information. Furthermore, in the treatment of degenerative conditions, compositions containing FGF-20 can be administered and it would likely be desirable to achieve certain target levels of FGF-20 in sera, in cerebrospinal fluid or in any desired tissue compartment. It would, therefore, be advantageous to be able to monitor the levels of FGF-20 in a patient. Accordingly, the present invention also provides methods for detecting the presence of FGF-20 in a sample from a patient.

The term "detection" as used herein in the context of detecting the presence of FGF-20 in a patient is intended to include the determining of the amount of FGF-20 or the ability to express an amount of FGF-20 in a patient, the distinguishing of FGF-20 from other growth factors, the estimation of prognosis in terms of probable outcome of a degenerative disease and prospect for recovery, the monitoring of the FGF-20 levels over a period of time as a measure of status of the condition, and the monitoring of FGF-20 levels for determining a preferred therapeutic regimen for the patient.

To detect the presence of FGF-20 in a patient, a sample is obtained from the patient. The sample can be a tissue biopsy sample or a sample of blood, plasma, serum,

CSF or the like. FGF-20 is expressed in neural tissues as discussed in Example 8. Samples for detecting FGF-20 can be taken from this tissue. When assessing peripheral levels of FGF-20, it is preferred that the sample be a sample of blood, plasma or serum. When assessing the levels of FGF-20 in the central nervous system a preferred sample is a sample
 5 obtained from cerebrospinal fluid or neural tissue.

In some instances it is desirable to determine whether the FGF-20 gene is intact in the patient or in a tissue or cell line within the patient. By an intact FGF-20 gene it is meant that there are no alterations in the gene such as point mutations, deletions, insertions, chromosomal breakage, chromosomal rearrangements and the like wherein such
 10 alteration might alter production of FGF-20 or alter its biological activity, stability or the like to lead to disease processes or susceptibility to cellular degenerative conditions. Thus, in one embodiment of the present invention a method is provided for detecting and characterizing any alterations in the FGF-20 gene. The method comprises providing an oligonucleotide that contains the FGF-20 cDNA, genomic DNA or a fragment thereof or a
 15 derivative thereof. By a derivative of an oligonucleotide, it is meant that the derived oligonucleotide is substantially the same as the sequence from which it is derived in that the derived sequence has sufficient sequence complementarity to the sequence from which it is derived to hybridize to the FGF-20 gene. The derived nucleotide sequence is not necessarily physically derived from the nucleotide sequence, but may be generated in any
 20 manner including for example, chemical synthesis or DNA replication or reverse transcription or transcription.

Typically, patient genomic DNA is isolated from a cell sample from the patient and digested with one or more restriction endonucleases such as, for example, TaqI and AluI. Using the Southern blot protocol, which is well known in the art, this assay
 25 determines whether a patient or a particular tissue in a patient has an intact FGF-20 gene or an FGF-20 gene abnormality.

Hybridization to an FGF-20 gene would involve denaturing the chromosomal DNA to obtain a single-stranded DNA; contacting the single-stranded DNA with a gene probe associated with the FGF-20 gene sequence; and identifying the

hybridized DNA-probe to detect chromosomal DNA containing at least a portion of a human FGF-20 gene.

The term “probe” as used herein refers to a structure comprised of a polynucleotide that forms a hybrid structure with a target sequence, due to complementarity of probe sequence with a sequence in the target region. Oligomers suitable for use as probes may contain a minimum of about 8-12 contiguous nucleotides which are complementary to the targeted sequence and preferably a minimum of about 20.

The FGF-20 gene probes of the present invention can be DNA or RNA oligonucleotides and can be made by any method known in the art such as, for example, excision, transcription or chemical synthesis. Probes may be labeled with any detectable label known in the art such as, for example, radioactive or fluorescent labels or enzymatic marker. Labeling of the probe can be accomplished by any method known in the art such as by PCR, random priming, end labeling, nick translation or the like. One skilled in the art will also recognize that other methods not employing a labeled probe can be used to determine the hybridization. Examples of methods that can be used for detecting hybridization include Southern blotting, fluorescence in situ hybridization, and single-strand conformation polymorphism with PCR amplification.

Hybridization is typically carried out at 25° - 45°C, more preferably at 32° - 40°C and more preferably at 37° - 38°C. The time required for hybridization is from about 0.25 to about 96 hours, more preferably from about one to about 72 hours, and most preferably from about 4 to about 24 hours.

FGF-20 gene abnormalities can also be detected by using the PCR method and primers that flank or lie within the FGF-20 gene. The PCR method is well known in the art. Briefly, this method is performed using two oligonucleotide primers which are capable of hybridizing to the nucleic acid sequences flanking a target sequence that lies within an FGF-20 gene and amplifying the target sequence. The terms “oligonucleotide primer” as used herein refers to a short strand of DNA or RNA ranging in length from about 8 to about 30 bases. The upstream and downstream primers are typically from about 20 to about 30 base pairs in length and hybridize to the flanking regions for replication of

the nucleotide sequence. The polymerization is catalyzed by a DNA-polymerase in the presence of deoxynucleotide triphosphates or nucleotide analogs to produce double-stranded DNA molecules. The double strands are then separated by any denaturing method including physical, chemical or enzymatic. Commonly, a method of physical denaturation
 5 is used involving heating the nucleic acid, typically to temperatures from about 80°C to 105°C for times ranging from about 1 to about 10 minutes. The process is repeated for the desired number of cycles.

The primers are selected to be substantially complementary to the strand of DNA being amplified. Therefore, the primers need not reflect the exact sequence of the
 10 template, but must be sufficiently complementary to selectively hybridize with the strand being amplified.

After PCR amplification, the DNA sequence comprising FGF-20 or pre-pro FGF-20 or a fragment thereof is then directly sequenced and analyzed by comparison of the sequence with the sequences disclosed herein to identify alterations which might change
 15 activity or expression levels or the like.

In another embodiment, a method for detecting FGF-20 is provided based upon an analysis of tissue expressing the FGF-20 gene, as described in the Examples. The method comprises hybridizing a polynucleotide to mRNA from a sample of tissue that normally expresses the FGF-20 gene. The sample is obtained from a patient suspected of
 20 having an abnormality in the FGF-20 gene or in the FGF-20 gene of particular cells.

To detect the presence of mRNA encoding FGF-20 protein, a sample is obtained from a patient. The sample can be from blood or from a tissue biopsy sample. The sample may be treated to extract the nucleic acids contained therein. The resulting nucleic acid from the sample is subjected to gel electrophoresis or other size separation
 25 techniques.

The mRNA of the sample is contacted with a DNA sequence serving as a probe to form hybrid duplexes. The use of a labeled probes as discussed above allows detection of the resulting duplex.

When using the cDNA encoding FGF-20 protein or a derivative of the cDNA as a probe, high stringency conditions can be used in order to prevent false positives, that is the hybridization and apparent detection of FGF-20 nucleotide sequences when in fact an intact and functioning FGF-20 gene is not present. When using sequences derived from the FGF-20 cDNA, less stringent conditions could be used, however, this would be a less preferred approach because of the likelihood of false positives. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, length of time and concentration of formamide.

In order to increase the sensitivity of the detection in a sample of mRNA encoding the FGF-20 protein, the technique of reverse transcription/polymerization chain reaction (RT/PCR) can be used to amplify cDNA transcribed from mRNA encoding the FGF-20 protein. The method of RT/PCR is well known in the art, and can be performed as follows. Total cellular RNA is isolated by, for example, the standard guanidium isothiocyanate method and the total RNA is reverse transcribed. The reverse transcription method involves synthesis of DNA on a template of RNA using a reverse transcriptase enzyme and a 3' end primer. Typically, the primer contains an oligo(dT) sequence. The cDNA thus produced is then amplified using the PCR method and FGF-20 specific primers. (Belyavsky et al., *Nucl. Acid Res.* 17:2919-2932 (1989); Krug and Berger, *Methods in Enzymology* 152:316-325, Academic Press, NY, 1987, which are incorporated by reference).

The polymerase chain reaction method is performed as described above using two oligonucleotide primers that are substantially complementary to the two flanking regions of the DNA segment to be amplified.

Following amplification, the PCR product is then electrophoresed and detected by ethidium bromide staining or by phosphoimaging.

The present invention further provides for methods to detect the presence of FGF-20 protein in a sample obtained from a patient. Any method known in the art for detecting proteins can be used. Such methods include, but are not limited to

immunodiffusion, immunoelectrophoresis, immunochemical methods, binder-ligand assays, immunohistochemical techniques, agglutination and complement assays. (For example, see *Basic and Clinical Immunology*, 217-262, Sites and Terr, eds., Appleton & Lange, Norwalk, CT, 1991 which is incorporated by reference). Preferred are binder-ligand

5 immunoassay methods including reacting antibodies with an epitope or epitopes of the FGF-20 protein and competitively displacing a labeled FGF-20 protein or derivative thereof. Preferred antibodies are prepared according to Example 11.

As used herein, a derivative of the FGF-20 protein is intended to include a polypeptide in which certain amino acids have been deleted, replaced, or changed to

10 modified or unusual amino acids wherein the FGF-20 derivative is biologically equivalent to FGF-20 and wherein the polypeptide derivative cross-reacts with antibodies raised against the FGF-20 protein. By cross-reaction it is meant that an antibody reacts with an antigen other than the one that induced its formation.

Numerous competitive and non-competitive protein binding immunoassays

15 are well known in the art. Antibodies employed in such assays may be unlabeled, for example as used in agglutination tests, or labeled for use in a wide variety of assay methods. Labels that can be used include radionuclides, enzymes, fluorescers, chemiluminescers, enzyme substrates or co-factors, enzyme inhibitors, particles, dyes and the like for use in radioimmunoassay (RIA), enzyme immunoassays, *e.g.*, enzyme-linked

20 immunosorbent assay (ELISA), fluorescent immunoassays and the like.

Polyclonal or monoclonal antibodies to the FGF-20 protein or an epitope thereof can be made for use in immunoassays by any of a number of methods known in the art. By epitope reference is made to an antigenic determinant of a polypeptide. An epitope could comprise 3 amino acids in a spatial conformation which is unique to the epitope.

25 Generally an epitope consists of at least 5 such amino acids. Methods of determining the spatial conformation of amino acids are known in the art, and include, for example, x-ray crystallography and 2 dimensional nuclear magnetic resonance.

One approach for preparing antibodies to a protein is the selection and preparation of an amino acid sequence of all or part of the protein, chemically synthesizing

the sequence and injecting it into an appropriate animal, usually a rabbit or a mouse (see Example 11).

Oligopeptides can be selected as candidates for the production of an antibody to the FGF-20 protein based upon the oligopeptides lying in hydrophilic regions, which are thus likely to be exposed in the mature protein. Oligopeptides for raising antibodies include the contiguous amino acids at positions 176-189, or 56-70 of SEQ ID NO:4. These oligopeptides are RDGARSKRHQKFTH (SEQ ID NO:5) and QLAHLHGILRRRQLY (SEQ ID NO:6). Additional oligopeptides can be determined using, for example, the Antigenicity Index of Welling, G.W. et al., *FEBS Lett.* 188:215-218 (1985), incorporated herein by reference.

Antibodies to FGF-20 can also be raised against oligopeptides that include one or more of the conserved regions identified herein such that the antibody can cross-react with other family members. Such antibodies can be used to identify and isolate the other family members.

Methods for preparation of the FGF-20 protein or an epitope thereof include, but are not limited to chemical synthesis, recombinant DNA techniques or isolation from biological samples. Chemical synthesis of a peptide can be performed, for example, by the classical Merrifield method of solid phase peptide synthesis (Merrifield, *J., Am. Chem. Soc.* 85:2149 (1963), which is incorporated by reference) or the FMOC strategy on a Rapid Automated Multiple Peptide Synthesis system (E.I. du Pont de Nemours Company, Wilmington, Delaware) (Caprino and Han, *J. Org. Chem.* 37:3404 (1972), which is incorporated by reference).

Polyclonal antibodies can be prepared by immunizing rabbits or other animals by injecting antigen followed by subsequent boosts at appropriate intervals. The animals are bled and sera assayed against purified FGF-20 protein usually by ELISA or by bioassay based upon the ability to block the action of FGF-20 on neurons or other cells. When using avian species, *e.g.*, chicken, turkey and the like, the antibody can be isolated from the yolk of the egg. Monoclonal antibodies can be prepared after the method of Milstein and Kohler by fusing splenocytes from immunized mice with continuously

replicating tumor cells such as myeloma or lymphoma cells. (Milstein and Kohler, *Nature* 256:495-497 (1975); Gutfre and Milstein, *Methods in Enzymology: Immunochemical Techniques* 73:1-46, Langone and Banatis eds., Academic Press, 1981 which are incorporated by reference). The hybridoma cells so formed are then cloned by limiting
 5 dilution methods and supernates assayed for antibody production by ELISA, RIA or bioassay.

The unique ability of antibodies to recognize and specifically bind to target proteins provides an approach for treating an overexpression of the protein. Thus, another aspect of the present invention provides for a method for preventing or treating diseases
 10 involving overexpression of the FGF-20 protein by treatment of a patient with specific antibodies to the FGF-20 protein.

Specific antibodies, either polyclonal or monoclonal, to the FGF-20 protein can be produced by any suitable method known in the art as discussed above. For example, murine or human monoclonal antibodies can be produced by hybridoma technology or,
 15 alternatively, the FGF-20 protein, or an immunologically active fragment thereof, or an anti-idiotypic antibody, or fragment thereof can be administered to an animal to elicit the production of antibodies capable of recognizing and binding to the FGF-20 protein. Such antibodies can be from any class of antibodies including, but not limited to IgG, IgA, IgM, IgD, and IgE or in the case of avian species, IgY and from any subclass of antibodies.

20 Polypeptides encoded by the instant polynucleotides and corresponding full-length genes can be used to screen peptide libraries, protein libraries, small molecule libraries, and phage display libraries, and other known methods, to identify analogs or antagonists.

Native FGF polypeptides may play a role in cancer. For example, FGF
 25 family members can induce marked morphological transformation of NIH 3T3 cells, and exhibit strong tumorigenicity in nude mice. Angiogenic activity has been exhibited by FGF family members. Thus, inhibitors of FGF can be used to treat cancer.

A library of peptides may be synthesized following the methods disclosed in U.S. Patent No. 5,010,175, and in PCT No. WO 91/17823. As described below in brief, a

mixture of peptides is prepared, which is then screened to identify the peptides exhibiting the desired signal transduction and receptor binding activity. According to the method of the '175 patent, a suitable peptide synthesis support (*e.g.*, a resin) is coupled to a mixture of appropriately protected, activated amino acids. The concentration of each amino acid in the reaction mixture is balanced or adjusted in inverse proportion to its coupling reaction rate so that the product is an equimolar mixture of amino acids coupled to the starting resin. The bound amino acids are then deprotected, and reacted with another balanced amino acid mixture to form an equimolar mixture of all possible dipeptides. This process is repeated until a mixture of peptides of the desired length (*e.g.*, hexamers) is formed. Note that one need not include all amino acids in each step: one may include only one or two amino acids in some steps (*e.g.*, where it is known that a particular amino acid is essential in a given position), thus reducing the complexity of the mixture. After the synthesis of the peptide library is completed, the mixture of peptides is screened for binding to the selected polypeptide. The peptides are then tested for their ability to inhibit or enhance activity. Peptides exhibiting the desired activity are then isolated and sequenced.

The method described in PCT No. WO 91/17823 is similar. However, instead of reacting the synthesis resin with a mixture of activated amino acids, the resin is divided into twenty equal portions (or into a number of portions corresponding to the number of different amino acids to be added in that step), and each amino acid is coupled individually to its portion of resin. The resin portions are then combined, mixed, and again divided into a number of equal portions for reaction with the second amino acid. In this manner, each reaction may be easily driven to completion. Additionally, one may maintain separate "subpools" by treating portions in parallel, rather than combining all resins at each step. This simplifies the process of determining which peptides are responsible for any observed receptor binding or signal transduction activity.

In such cases, the subpools containing, *e.g.*, 1-2,000 candidates each are exposed to one or more polypeptides of the invention. Each subpool that produces a positive result is then resynthesized as a group of smaller subpools (sub-subpools) containing, *e.g.*, 20-100 candidates, and reassayed. Positive sub-subpools may be

resynthesized as individual compounds, and assayed finally to determine the peptides that exhibit a high binding constant. These peptides can be tested for their ability to inhibit or enhance the native activity. The methods described in PCT No. WO 91/7823 and U.S. Patent No. 5,194,392 (herein incorporated by reference) enable the preparation of such pools and subpools by automated techniques in parallel, such that all synthesis and resynthesis may be performed in a matter of days.

Peptide agonists or antagonists are screened using any available method, such as signal transduction, antibody binding, receptor binding and mitogenic assays. The assay conditions ideally should resemble the conditions under which the native activity is exhibited *in vivo*, that is, under physiologic pH, temperature, and ionic strength. Suitable agonists or antagonists will exhibit strong inhibition or enhancement of the native activity at concentrations that do not cause toxic side effects in the subject. Agonists or antagonists that compete for binding to the native polypeptide may require concentrations equal to or greater than the native concentration, while inhibitors capable of binding irreversibly to the polypeptide may be added in concentrations on the order of the native concentration.

The availability of hFGF-20 and rFGF-20 allows for the identification of small molecules and low molecular weight compounds that inhibit the binding of FGF-20 to its receptor, through routine application of high-throughput screening methods (HTS). HTS methods generally refer to technologies that permit the rapid assaying of lead compounds for therapeutic potential. HTS techniques employ robotic handling of test materials, detection of positive signals, and interpretation of data. Lead compounds may be identified via the incorporation of radioactivity or through optical assays that rely on absorbance, fluorescence or luminescence as read-outs. Gonzalez, J.E., et al., *Curr. Opin. Biotech.* 9:624-631 (1998). Assays for detecting interaction between an FGF molecule and FGF receptor are described in, for example, Blunt, A. G. et al., *J. Biol. Chem.* 272:3733-3738 (1997), and such assays can be adapted for determining if a candidate molecule can inhibit the interaction between FGF-20 and its receptor.

Model systems are available that can be adapted for use in high throughput screening for compounds that inhibit the interaction of FGF-20 with receptors to which it

binds (see Example 12), for example by competing with FGF-20 for receptor binding. Sarubbi et al., *Anal. Biochem.* 237:70-75 (1996), describe cell-free, non-isotopic assays for discovering molecules that compete with natural ligands for binding to the active site of IL-1 receptor. Martens, C. et al., *Anal. Biochem.* 273:20-31 (1999), describe a generic particle-based nonradioactive method in which a labeled ligand binds to its receptor immobilized on a particle; label on the particle decreases in the presence of a molecule that competes with the labeled ligand for receptor binding.

The therapeutic FGF-20 polynucleotides and polypeptides of the present invention may be utilized in gene delivery vehicles. The gene delivery vehicle may be of viral or non-viral origin (*see generally*, Jolly, *Cancer Gene Therapy* 1:51-64 (1994); Kimura, *Human Gene Therapy* 5:845-852 (1994); Connelly, *Human Gene Therapy* 1:185-193 (1995); and Kaplitt, *Nature Genetics* 6:148-153 (1994)). Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches. Expression of such coding sequences can be induced using endogenous mammalian or heterologous promoters. Expression of the coding sequence can be either constitutive or regulated.

The present invention can employ recombinant retroviruses which are constructed to carry or express a selected nucleic acid molecule of interest. Retrovirus vectors that can be employed include those described in EP 0 415 731; WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 93/11230; WO 93/10218; Vile and Hart, *Cancer Res.* 53:3860-3864 (1993); Vile and Hart, *Cancer Res.* 53:962-967 (1993); Ram et al., *Cancer Res.* 53:83-88 (1993); Takamiya et al., *J. Neurosci. Res.* 33:493-503 (1992); Baba et al., *J. Neurosurg.* 79:729-735 (1993); U.S. Patent No. 4,777,127; GB Patent No. 2,200,651; and EP 0 345 242. Preferred recombinant retroviruses include those described in WO 91/02805.

Packaging cell lines suitable for use with the above-described retroviral vector constructs may be readily prepared (*see* PCT publications WO 95/30763 and WO 92/05266), and used to create producer cell lines (also termed vector cell lines) for the

production of recombinant vector particles. Within particularly preferred embodiments of the invention, packaging cell lines are made from human (such as HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviruses that can survive inactivation in human serum.

5 The present invention also employs alphavirus-based vectors that can function as gene delivery vehicles. Such vectors can be constructed from a wide variety of alphaviruses, including, for example, Sindbis virus vectors, Semliki forest virus (ATCC VR-67; ATCC VR-1247), Ross River virus (ATCC VR-373; ATCC VR-1246) and Venezuelan equine encephalitis virus (ATCC VR-923; ATCC VR-1250; ATCC VR 1249; 10 ATCC VR-532). Representative examples of such vector systems include those described in U.S. Patent Nos. 5,091,309; 5,217,879; and 5,185,440; and PCT Publication Nos. WO 92/10578; WO 94/21792; WO 95/27069; WO 95/27044; and WO 95/07994.

Gene delivery vehicles of the present invention can also employ parvovirus such as adeno-associated virus (AAV) vectors. Representative examples include the AAV 15 vectors disclosed by Srivastava in WO 93/09239, Samulski et al., *J. Vir.* 63:3822-3828 (1989); Mendelson et al., *Virol.* 166:154-165 (1988); and Flotte et al., *P.N.A.S.* 90:10613-10617 (1993).

AAV vectors may be suitable for administering FGF-20 to treat hearing disorders. For example, Lalwani et al., *Gene Ther.* 3:588-592 (1996), used AAV to obtain 20 *in vivo* expression of a foreign gene in the cochlea of guinea pigs.

Representative examples of adenoviral vectors include those described by Berkner, *Biotechniques* 6:616-627 (Biotechniques); Rosenfeld et al., *Science* 252:431-434 (1991); WO 93/19191; Kolls et al., *P.N.A.S.* :215-219 (1994); Kass-Eisler et al., *P.N.A.S.* 90:11498-11502 (1993); Guzman et al., *Circulation* 88:2838-2848 (1993); Guzman et al., 25 *Cir. Res.* 73:1202-1207 (1993); Zabner et al., *Cell* 75:207-216 (1993); Li et al., *Hum. Gene Ther.* 4:403-409 (1993); Cailaud et al., *Eur. J. Neurosci.* 5:1287-1291 (1993); Vincent et al., *Nat. Genet.* 5:130-134 (1993); Jaffe et al., *Nat. Genet.* 1:372-378 (1992); and Levrero et al., *Gene* 101:195-202 (1992). Exemplary adenoviral gene therapy vectors employable in this invention also include those described in WO 94/12649, WO 93/03769; WO 93/19191;

WO 94/28938; WO 95/11984 and WO 95/00655. Administration of DNA linked to killed adenovirus as described in Curiel, *Hum. Gene Ther.* 3:147-154 (1992), may be employed.

Other gene delivery vehicles and methods may be employed, including polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example Curiel, *Hum. Gene Ther.* 3:147-154 (1992); ligand-linked DNA, for example see Wu, *J. Biol. Chem.* 264:16985-16987 (1989); eukaryotic cell delivery vehicles cells, for example see U.S. Serial No. 08/240,030, filed May 9, 1994, and U.S. Serial No. 08/404,796; deposition of photopolymerized hydrogel materials; hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; ionizing radiation as described in U.S. Patent No. 5,206,152 and in WO 92/11033; nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip, *Mol. Cell Biol.* 14:2411-2418 (1994), and in Woffendin, *Proc. Natl. Acad. Sci.* 91:1581-1585 (1994).

Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and U.S. Patent No. 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm. Liposomes that can act as gene delivery vehicles are described in U.S. Patent No. 5,422,120, PCT Patent Publication Nos. WO 95/13796, WO 94/23697, and WO 91/14445, and EP No. 0 524 968.

Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin et al., *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585 (1994). Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; use of ionizing radiation for activating transferred gene, as described in U.S. Patent No. 5,206,152 and PCT Patent Publication No. WO 92/11033.

Loss of pulmonary, bronchia or alveolar cells or function, healing of pulmonary or bronchia wounds, pulmonary infraction, emphysema/chronic obstructive pulmonary disease, asthma, sequelae of infectious or autoimmune disease, sequelae of pulmonary arterial or venous hypertension, pulmonary fibrosis, pulmonary disease of immaturity, and cystic fibrosis are conditions amenable to treatment with FGF.

FGF-20 polynucleotides and polypeptides may aid in correction of congenital defects due to loss of FGF-20 molecule or its function (heart, lung, brain, limbs, kidney, etc.). FGF-20 polynucleotides and polypeptides may also aid in the correction of such defects wherein the defects lead to hearing loss due to cochlear defects.

40

FGF-20 may also be used in screens to identify drugs for treatment of cancers which involve over activity of the molecule, or new targets which would be useful in the identification of new drugs.

For all of the preceding embodiments, the clinician will determine, based on the specific condition, whether FGF-20 polypeptides or polynucleotides, antibodies to FGF-20, or small molecules such as peptide analogues or antagonists, will be the most suitable form of treatment. These forms are all within the scope of the invention.

Preferred embodiments of the invention are described in the following examples. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

15 EXAMPLE 1

20 EXAMPLE 2

41

rat FGF-9 (17) and FGF-16 (21), FEENWY and THFLPR, respectively. The amplified product was further amplified by PCR with each of the sense and antisense degenerate primers representing all possible codons corresponding to another consensus amino acid sequences of rat FGF-9 (17) and FGF-16 (21), ENWYNT and HQKFTH, respectively. The amplified DNA of expected size (approximately 150 base pairs) was cloned into the pGEM-T DNA vector (Promega, Madison, Wisconsin). The nucleotide sequence of the cloned DNA was determined by a DNA sequencer (Applied Biosystems, Foster, California). To determine the coding region of a novel FGF cDNA, the coding region was amplified from cDNA synthesized from rat brain poly (A)⁺ RNA by adaptor-ligation mediated polymerase chain reaction using a Marathon cDNA amplification kit (Clontech, Palo Alto, California). To determine the amino-terminal region, DNA encoding the region was amplified from rat genomic DNA by cassette-ligation mediated polymerase chain reaction (Isegawa, Y. et al., *Mol. Cell. Probes* 6:467-475 (1992)) using a LA PCR *in vitro* cloning kit (TaKaRa, Kyoto, Japan). The cDNA encoding the entire coding region of the FGF was amplified from rat brain cDNA by polymerase chain reaction in the presence of 5% dimethyl sulfoxide (Villarreal, X.C. et al., *Anal. Biochem.* 197:362-367 (1991)) using the FGF-specific primers including the 5' and 3' noncoding sequences, and cloned into the pGEM-T DNA vector. The apparent evolutionary relationships of members of the FGF family were examined by the unweighted pair-group method with arithmetic mean method using the sequence analysis software, Genetyx (Software Development Co., Tokyo, Japan).

EXAMPLE 3

Expression of FGF-20 mRNA in Rat Embryonic Inner Ear --- Expression of FGF-20 in rat embryos was examined. Consecutive transverse sections of rat embryos (E14.5) were examined by *in situ* hybridization with ³⁵S-labeled FGF-20 anti-sense and sense cRNA probes. The sections were counterstained with hematoxylin and eosin. Bright-field and dark-field photographs of the sections reveal that FGF-20 is preferentially expressed in the cochlea of the inner ear.

EXAMPLE 4

Northern Blotting Analysis --- Poly (A)⁺ RNA (10 µg) from rat adult brain was dissolved on a denaturing agarose gel (1%) containing formaldehyde, and transferred to a nitrocellulose membrane in 20X SSC (1X SSC:0.15 M NaCl/0.015 M sodium citrate) overnight. A ³²P-labeled FGF-20 cDNA probe (~650 base pairs) was labeled with a random primer labeling kit (Pharmacia Biotech, Uppsala, Sweden) and deoxycytidine 5'-[α-³²P-] triphosphate (~110 TBq/mmol) (ICN Biomedicals Inc., Costa Mesa, California). The membrane was incubated in hybridization solution containing the labeled probe as described (22), and analyzed with a radio-imaging analyzer (BAS 2000, Fuji Photo Film Co., Tokyo, Japan). To confirm the integrity of the poly (A)⁺ RNA, the hybridized probe on the membrane was washed with 0.5X SSC containing 0.01 M EDTA (pH 8.0) at 100°C for 5 min and with 0.05X SSC containing 0.01 M EDTA (pH 8.0) and 0.1% SDS at 60°C for 15 min. The washed membrane was rehybridized with a ³²P-labeled rat βactin cDNA probe (~410 base pairs) (Nudel, U. et al., *Nucleic Acids Res.* 11:1759-1771 (1983)).

15 EXAMPLE 5

In Situ Hybridization --- Adult Wistar rat brain was frozen in powdered dry ice, and sagittal sections were cut at 16 µm with a cryostat, thaw-mounted onto poly-L-lysine-coated slides, and stored at -85°C until hybridization. A ³⁵S-labeled rat FGF-20 antisense or sense cRNA probe was transcribed using SP6 RNA polymerase or T7 RNA polymerase (TaKaRa, Kyoto, Japan) with uridine 5'-α[³⁵S]thiotriphosphate (~30 TBq/mmol) (Amersham, Buckinghamshire, England), respectively. The sections were examined by *in situ* hybridization with the labeled probe as described (Yamasaki, M. et al., *J. Biol. Chem.* 271:15918-15921 (1996)).

EXAMPLE 6

Preparation of Recombinant Rat FGF-20 --- The rat FGF-20 cDNA with a DNA fragment (75 BP) encoding an E-tag (GAPVPYPDPLEPR) and a His₆ tag (HHHHHH) at the 3'-terminus of the coding region was constructed in a transfer vector DNA, *pBacPAK9* (Clontech, Palo Alto, California). Recombinant baculovirus containing the FGF-20 cDNA with the tag sequences was obtained by cotransfection of Sf9 cells with the recombinant *pBacPAK9* and a Bsu36 I-digested expression vector, *BacPAK6* (Clontech, Palo Alto, California). High Five insect cells were infected with the resultant recombinant baculovirus and incubated at 27°C for 65 h in serum-free medium EX-CELL 400 (JRH Biosciences, Lenexa, Kansas). The culture medium was dialyzed against phosphate-buffered saline (PBS), and applied to a column of Ni-NTA agarose (QIAGEN GmbH, Hilden, Germany) in PBS containing 20 mM imidazole and 0.5 M NaCl. After washing the column with PBS containing 20 mM imidazole and 0.5 M NaCl, recombinant FGF-20 was eluted from the column with PBS containing 250 mM imidazole and 0.5 M NaCl, and applied to a column of Bio-Gel P-6 DG (Bio-Rad Lab., Hercules, California) in PBS containing 100 μ g/ml BSA.

EXAMPLE 7

Detection of Recombinant FGF-20 by Western Blotting Analysis --- The culture medium or rat recombinant FGF-20 was separated by sodium dodecyl sulfate (SDS)-polyacrylamide gel (12.5%) electrophoresis under reducing conditions and transferred onto a nitrocellulose membrane (Hybond-ECL, Amersham, Buckinghamshire, England). The membrane was incubated with anti-E tag antibodies (1:500) (Pharmacia Biotech, Uppsala, Sweden). The protein with the E-tag was visualized as described (Hoshikawa, M. et al., *Biochem. Biophys. Res. Commun.* 244:187-191 (1998)).

EXAMPLE 8

Rat Midbrain Cultured Cells --- The ventral mesencephalon was resected from rat embryos (E16.5). The mesencephalic blocks were washed 10 times with Hanks' solution and mechanically dissociated without enzymatic treatment. The midbrain cultured cells were prepared essentially as described (Sawada, H. et al., *J. Neurosci. Res.* 43:503-510 (1996)). The culture medium consisted of Eagle's minimum essential medium (EMEM) supplemented with 0.2% sodium carbonate, 0.1% glucose, 0.029% L-glutamine and 0.238% HEPES. The cultured cells were incubated at 37°C in the culture medium containing 10% fetal calf serum. From the 5th day of culture, the cells were incubated in the culture medium containing 10% horse serum.

EXAMPLE 9

Examination of Neurotrophic Activity of FGF-20 for Midbrain Dopaminergic Neurons --- Cells on the 8th day of culture were incubated in Eagle's minimum essential medium supplemented with 0.2% sodium hydrogen carbonate, 0.1% glucose, 0.029% L-glutamine, 0.238% HEPES and 10% horse serum or 0.1% bovine serum albumin in the presence or absence of FGF-20 for 4 days and then fixed with fresh 4% paraformaldehyde for 30 min on the 12th day. Cells on the 8th day of culture were also incubated in the presence or absence of recombinant rat FGF-20 for 24 h and then were treated with 1 mM glutamate for 10 min. The cultured cells were further incubated in medium without FGF-20 and 1mM glutamate and then fixed with 4% paraformaldehyde for 30 min on the 12th day. The fixed cells were washed with PBS for 15 min, and then treated with 0.2% Triton X-10 for 30 min. The cells were immunostained with anti-tyrosine hydroxylase (TH) antibody (Eugene Tech, Ridgefield Park, New Jersey) essentially as described (Sawada, H. et al., *J. Neurosci. Res.* 43:503-510 (1996)). Numbers of cultured dopaminergic neurons were evaluated by counting cells stained with anti-TH antibody.

EXAMPLE 10

Isolation and Analysis of Human FGF-20 --- The coding region of human FGF-20 DNA was amplified from human brain cDNA library (λ gt10) by PCR using primers specific for FGF-20 and λ gt10 DNA. The nucleotide sequence of the cDNA encoding the carboxy-terminal 112 amino acids of human FGF-20 was determined, and is shown in Figure 7. An alignment of rat and human FGF-20 amino acid sequences is shown in Figure 8.

EXAMPLE 11

Preparation of Antisera to FGF-20 by Immunization of Rabbits with an FGF-20 Peptide --- A peptide sequence corresponding to selected contiguous amino acids of the human FGF-20 protein is synthesized and coupled to keyhole limpet hemocyanin (KLH) as described (Harlow and Land, Antibodies: A Laboratory Manual, 1988. Cold Spring Harbor Laboratory, New York, NY). The KLH-coupled peptide is used to immunize rabbits. Antisera are tested for specificity to FGF-20, and for cross-reactivity with other FGF proteins.

Exemplary peptide sequences are:

1. RDGARSKRHQKFTH (SEQ ID NO:5)
2. QLAHLHGILRRRQLY (SEQ ID NO:6)

EXAMPLE 12

Binding of FGF-20 to the Recombinant Extracellular Domains of FGFR-1c, FGFR-2c, and FGFR-3c -- Recombinant FGF-20 was fixed on the sensor tip CM5 (Amersham Pharmacia Biotech). Binding of the recombinant extracellular domain of FGFR-1c, FGFR-2c, or FGFR-3c to FGF-20 on the tip was analyzed using the BIACORE

2000 System (Amersham Pharmacia Biotech). The equilibrium dissociation constant was determined by the BIA evaluation software (Amersham Pharmacia Biotech).

TABLE 1

5

BINDING OF FGF-20 TO FGF RECEPTORS

Receptor	K_{diss} (S^{-1})	K_{ass} ($\text{M}^{-1} \bullet \text{S}^{-1}$)	K_{d} (M)
FGFR-1c	nd	nd	
FGFR-2c	1.67×10^{-2}	5.95×10^5	2.81×10^{-8}
FGFR-3c	2.47×10^{-2}	1.15×10^5	2.17×10^{-7}

nd: not detected

As shown in Table 1, FGF-20 binds to FGF receptors 2 and 3, but not to FGF receptor 1. Thus, FGF-20 may exhibit biological effects not found in members of the FGF family that bind to FGF receptor 1, such as FGF-2 and FGF-4.

10

All patents, published patent applications and publications cited herein are incorporated by reference as if set forth fully herein.

Although certain preferred embodiments have been described herein, it is not intended that such embodiments be construed as limitations on the scope of the invention except as set forth in the following claims.

CLAIMS

WE CLAIM:

1. An isolated nucleic acid molecule comprising a polynucleotide selected from the group consisting of:

(a) a polynucleotide encoding amino acids from about 1 to about 211 of SEQ ID NO:4;

(b) a polynucleotide encoding amino acids from about 2 to about 211 of SEQ ID NO:4;

(c) a polynucleotide encoding amino acids from about 1 to about 169 of SEQ ID NO:4;

(d) a polynucleotide encoding amino acids from about 187 to about 211 of SEQ ID NO:4;

(e) a polynucleotide encoding amino acids from about 170 to about 186 of SEQ ID NO:4;

(f) a polynucleotide encoding amino acids from about 59 to about 193 of SEQ ID NO:4;

(g) a polynucleotide encoding at least 30 contiguous amino acids of SEQ ID NO:4;

(h) a polynucleotide encoding the amino acid sequence of a fragment of the polypeptide shown as amino acids 1 to 211 of SEQ ID NO:4, wherein said fragment has fibroblast growth factor activity;

(i) the polynucleotide complement of (a), (b), (c), (d), (e), (f), (g), or (h); and

(j) a polynucleotide at least 90% identical to the polynucleotide of (a), (b), (c), (d), (e), (f), (g), or (h).

2. An isolated nucleic acid molecule which comprises 20-600 contiguous nucleotides from the coding region of SEQ ID NO:3.

3. The isolated nucleic acid molecule of claim 2, which comprises 60-400 contiguous nucleotides from the coding region of SEQ ID NO:3

4. The isolated nucleic acid molecule of claim 3, which comprises 200-300 contiguous nucleotides from the coding region of SEQ ID NO:3.

5. An isolated nucleic acid molecule comprising a polynucleotide encoding a polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has an amino acid sequence selected from the group consisting of:

- (a) amino acids from about 1 to about 211 of SEQ ID NO:4;
- (b) amino acids from about 2 to about 211 of SEQ ID NO:4;
- (c) amino acids from about 170 to about 186 of SEQ ID NO:4;
- (d) amino acids from about 1 to about 169 and from about 187 to about 211 of SEQ ID NO:4, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and
- (e) amino acids from about 59 to about 193 of SEQ ID NO:4.

6. The isolated nucleic acid molecule of claim 1, which is DNA.

7. A method of making a recombinant vector comprising inserting a nucleic acid molecule of claim 1 into a vector in operable linkage to a promoter.

8. A recombinant vector produced by the method of claim 7.

9. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 8 into a host cell.

10. A recombinant host cell produced by the method of claim 9.

12. An isolated polypeptide comprising amino acids at least 95% identical to amino acids selected from the group consisting of:

- SEQ ID NO:4, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

13. An isolated polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has an amino acid sequence selected from the group consisting of:

- SEQ ID NO:4, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

14. An isolated polypeptide comprising amino acids selected from the group consisting of:

- 50

- (c) amino acids from about 170 to about 186 of SEQ ID NO:4;
 - (d) amino acids from about 1 to about 169 and from about 187 to about 211 of SEQ ID NO:4, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and
 - (e) amino acids from about 59 to about 193 of SEQ ID NO:4.
15. An epitope-bearing portion of the polypeptide of SEQ ID NO:4.
 16. The epitope-bearing portion of claim 15, which comprises between 10 and 50 contiguous amino acids of SEQ ID NO:4.
 17. The epitope-bearing portion of claim 15, which comprises a polypeptide selected from the group consisting of amino acids RDGARSKRHQKFTH (SEQ ID NO:5) and QLAHLHGILRRRQLY (SEQ ID NO:6).
 18. An isolated antibody that binds specifically to the polypeptide of claim 12.
 19. An isolated antibody that binds specifically to the polypeptide of claim 13.
 20. An isolated antibody that binds specifically to the polypeptide of claim 14.
 21. A pharmaceutical composition comprising the polypeptide of claim 12, in combination with a pharmaceutically acceptable carrier.
 22. A method for providing trophic support for cells in a patient in need thereof, the method comprising administering to the patient a composition comprising a polynucleotide encoding the polypeptide of SEQ ID NO:4.

23. The method of claim 22 wherein the patient suffers from Parkinson's disease.

24. A method for providing trophic support for cells in a patient in need thereof, the method comprising administering to the patient a composition comprising a polypeptide of SEQ ID NO:4.

25. The method of claim 24 wherein the patient suffers from Parkinson's disease.

26. The method of claim 25 wherein the patient suffers from a condition affecting the substantia nigra.

27. A method of alleviating a disease condition in the brain of a human patient wherein said disease condition is alleviated by at least one method selected from the group consisting of slowing degeneration of, restoring function of, and increasing the number of, dopaminergic neurons in said human patient, said method comprising administering to said patient a pharmaceutically effective composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:4.

28. A method of alleviating a disease condition in the brain of a human patient wherein said disease condition is alleviated by at least one method selected from the group consisting of slowing degeneration of, restoring function of, and increasing the number of, dopaminergic neurons in said human patient, said method comprising administering to said patient a pharmaceutically effective composition comprising a polynucleotide encoding the amino acid sequence of SEQ ID NO:4.

29. The method of claim 28 wherein said polynucleotide has the sequence of SEQ ID NO:3.

30. A kit for detecting the presence of mRNA encoding FGF-20 in a sample from a patient, said kit comprising a polynucleotide having at least 20 contiguous nucleotides of the nucleic acid of claim 1, packaged in a container.

31. A kit for detecting the presence of FGF-20 polypeptide in a sample from a patient, said kit comprising an antibody according to claim 18, packaged in a container.

32. A method of alleviating a cochlea-associated disease condition in a human patient wherein said disease condition is alleviated by at least one method selected from the group consisting of slowing degeneration of, restoring normal structure of, and maintaining normal structure of, the cochlea in said human patient, said method comprising administering to said patient a pharmaceutically effective composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:4 or a polynucleotide encoding the amino acid sequence of SEQ ID NO:4.

33. The method of claim 32, wherein said disease condition is selected from the group consisting of otosclerosis; Cogan's syndrome; Meniere's disease; Pendred's syndrome; diabetes-associated hearing loss; congenital malformations; autoimmune disease-related hearing loss; age-related hearing loss; deafness associated with lack of FGF receptor; and ischemia-related hearing disturbance.

34. An isolated nucleic acid molecule comprising a polynucleotide selected from the group consisting of:

- (a) a polynucleotide encoding amino acids from about 1 to about 212 of SEQ ID NO:2;
- (b) a polynucleotide encoding amino acids from about 2 to about 212 of SEQ ID NO:2;
- (c) a polynucleotide encoding amino acids from about 1 to about 169 of SEQ ID NO:2;

(d) a polynucleotide encoding amino acids from about 187 to about 212 of SEQ ID NO:2;

(e) a polynucleotide encoding amino acids from about 170 to about 186 of SEQ ID NO:2;

(f) a polynucleotide encoding amino acids from about 59 to about 193 of SEQ ID NO:2;

(g) a polynucleotide encoding at least 30 contiguous amino acids of SEQ ID NO:2;

(h) a polynucleotide encoding the amino acid sequence of a fragment of the polypeptide shown as amino acids 1 to 212 of SEQ ID NO:2, wherein said fragment has fibroblast growth factor activity;

(i) a polynucleotide encoding the amino acid sequence of a fragment of the polypeptide shown as amino acids 1 to 212 of SEQ ID NO:2, wherein said fragment has fibroblast growth factor activity.

(j) the polynucleotide complement of (a), (b), (c), (d), (e), (f), (g), (h), or (i);
and

(k) a polynucleotide at least 90% identical to the polynucleotide of (a), (b), (c), (d), (e), (f), (g), (h), or (i).

35. An isolated nucleic acid molecule which comprises 20-600 contiguous nucleotides from the coding region of SEQ ID NO:1.

36. The isolated nucleic acid molecule of claim 35, which comprises 60-400 contiguous nucleotides from the coding region of SEQ ID NO:1.

37. The isolated nucleic acid molecule of claim 36, which comprises 200-300 contiguous nucleotides from the coding region of SEQ ID NO:1.

- (a) amino acids from about 1 to about 212 of SEQ ID NO:2;
- (b) amino acids from about 2 to about 212 of SEQ ID NO:2;
- (c) amino acids from about 170 to about 186 of SEQ ID NO:2;
- (d) amino acids from about 1 to about 169 and from about 187 to about 212 of SEQ ID NO:2, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

- (e) amino acids from about 59 to about 193 of SEQ ID NO:4.

39. The isolated nucleic acid molecule of claim 34, which is DNA.

40. A method of making a recombinant vector comprising inserting a nucleic acid molecule of claim 34 into a vector in operable linkage to a promoter.

41. A recombinant vector produced by the method of claim 40.

42. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 41 into a host cell.

43. A recombinant host cell produced by the method of claim 42.

44. A recombinant method of producing a polypeptide, comprising culturing the recombinant host cell of claim 43 under conditions such that said polypeptide is expressed and recovering said polypeptide.

45. An isolated polypeptide comprising amino acids at least 95% identical to amino acids selected from the group consisting of:

- (a) amino acids from about 1 to about 212 of SEQ ID NO:2;
- (b) amino acids from about 2 to about 212 of SEQ ID NO:2;
- (c) amino acids from about 170 to about 186 of SEQ ID NO:2;
- (d) amino acids from about 1 to about 169 and from about 187 to about 212 of

SEQ ID NO:2, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

- (e) amino acids from about 59 to about 193 of SEQ ID NO:2.

46. An isolated polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has an amino acid sequence selected from the group consisting of:

- (a) amino acids from about 1 to about 212 of SEQ ID NO:2;
- (b) amino acids from about 2 to about 212 of SEQ ID NO:2;
- (c) amino acids from about 170 to about 186 of SEQ ID NO:2;
- (d) amino acids from about 1 to about 169 and from about 187 to about 212 of

SEQ ID NO:2, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

- (e) amino acids from about 59 to about 193 of SEQ ID NO:2.

47. An isolated polypeptide comprising amino acids selected from the group consisting of:

- (a) amino acids from about 1 to about 212 of SEQ ID NO:2;
- (b) amino acids from about 2 to about 212 of SEQ ID NO:2;
- (c) amino acids from about 170 to about 186 of SEQ ID NO:2;
- (d) amino acids from about 1 to about 169 and from about 187 to about 212 of

SEQ ID NO:2, wherein said amino acids at positions about 169 and about 187 are joined by a peptide bond; and

- (e) amino acids from about 59 to about 193 of SEQ ID NO:2.
48. An epitope-bearing portion of the polypeptide of SEQ ID NO:2.
49. The epitope-bearing portion of claim 48, which comprises between 10 and 50 contiguous amino acids of SEQ ID NO:2.
50. An isolated antibody that binds specifically to the polypeptide of claim 45.
51. An isolated antibody that binds specifically to the polypeptide of claim 46.
52. An isolated antibody that binds specifically to the polypeptide of claim 47.

HUMAN FGF-20 GENE AND GENE EXPRESSION PRODUCTS

ABSTRACT OF THE DISCLOSURE


This invention relates to human fibroblast growth factor (hFGF-20), and to variants thereof and to polynucleotides encoding FGF-20. The invention also relates to diagnostic and therapeutic agents related to the polynucleotides and proteins, including probes and antibodies, to methods of treating neuronal degenerative disease such as Parkinson's disease and to methods of treating disorders of the cochlea including those causing hearing loss. The invention also relates to rat fibroblast growth factor (rFGF-20), and to variants thereof and polynucleotides encoding rFGF-20.

L \201130 - Chron\407\407-AP doc\4

Figure 1


rat FGF-9	MAPLGEVGSYF-GVCI)-AVPFGNPVL--PVDSPVLLSDHLGQSEAGGLPRGPAVTDLHL	57
	**** *	
rat FGF-20	MAPLTEVGAFLLGLEGLGQVGSFLLPPAGERPPLLGERRGALE-RGARGPGSGVELAHL	60
	*** *	
rat FGF-16	MAEVGGVFASLDWDLQGFSSSLGNVPLADSPGFLNERLQGIE--GKLQSGSPTDFAHL	96
	KGILRRRQLYCRTGFLEIFPNGTIQTRKDHRSFGILEFISIAVGLVSIRGVDSGLYLG	117
	***** *	
	HGILRRRQLYCRTGFLLQILPDGSVQGTQDHSFGILEFISVAVGLVSIRGVDSGLYLG	120
	***** *	
	KGILRRRQLYCRTGFHLEIFPNGTVHGRHDSRFGILEFISLAVGLISIRGVDSGLYLG	116
	MNEKGELYGSEKLTQECVFREQFEENWYNTYSSNL YKHVDTGRRYYVALNKDGTREGTR	177
	** ***** *	
	MNGKGELYGSEKLTSCIFREQFEENWYNTYSSNIYKHGDTGRRYFVALNKDGTPRDGR	180
	** ***** *	
	MNERGELFGSKKLTRICVFREQFEENWYNTYASTLYKHSDSERQYYVALNKDGSPPREGYR	176
	TKRHQKFTHFLPRPVIPDKVPPELYKDILSQS	208
	***** *	
	SKRHQKFTHFLPRPVIPERVPELYKDLLVYTG	212
	***** *	
	TKRHQKFTHFLPRVPDPSKLPMSRDLFRYR	207

Figure 2



000001 54526960

Figure 3



09692945-102000

Figure 4





Figure 5



09692945 102000

Figure 6

SEQ ID NO:1

Rat FGF-20 DNA

Translation start and stop codons indicated in bold, underlined type.

CCTTCC**ATG**GCTCCCTTGACCGAAGTCGGTGCCTTCTTGGGCGGCCTGGAGGGCTTGGGCCAGCAGGTGGGG
TCGCACTTCTTGCTGCCTCCTGCAGGGGAGCGACCGCCGCTGCTAGGGGAGCGGCGGGGCGCGTTGGAGCGG
GGCGCCCGCGGCGGGCCGGGTTCGGTGGAGCTGGCGCACCTGCACGGCATCCTGCGCCGCCGGCAGCTCTAC
TGCCGCACCGGCTTCCACCTGCAGATCCTGCCCCACGGCAGTGTGCAGGGCACCCGGCAGGATCACAGCCTC
TTCGGTATCCTGGAATTCATCAGTGTGGCGGTGGGGCTGGTCAGTATCAGAGGTGTGGACAGCGGCCTGTAC
CTTGGCATGAATGGCAAAGGAGAGCTTTATGGCTCAGAGAAATTGACTTCTGAATGCATCTTCAGGGAACAA
TTTGAAGAGAACTGGTATAATACCTATTCATCCAACATATACAAACACGGAGACACAGGTTCGCAGGTATTTT
GTAGCACTTAACAAAGACGGGACTCCAAGGGACGGTGCCAGGTCCAAAAGACACCAAAGTTTACCCATTTT
TTACCCAGACCAGTGGACCCAGAGAGAGTCCCAGAGTTATACAAAGACCTACTGGTGTACACTGGAT**TGA**ACC

000207" 54526960

Figure 7

SEQ ID NO:2

Amino acid sequence of rat FGF-20

MAPLTEVGAFLGGLEGLGQQVGSFLLPPAGERPPLLGERRGALERGARGGPGSVELAHLHGILRRRQLYCR
TGFHLQILPDGSVQGTRQDHSLEFGILEFISVAVGLVSIRGVDSGLYLG MNGKGELYGSEKLTSECIFREQFE
ENWYNTYSSNIYKHGDTGRRYFVALNKDGTPRDGARSKRHQKFTHFLRPVDPERVPELYKDLLVYTG

000207" 34626960

Figure 8

Sequence Size : 636 humanFGF-20 DNA (coding)

10 20 30 40 50 60
atggtcccttagccgaagtcgggggctttctgggcggcctggagggcttgggccagcag
M A P L A E V G G F L G G L E G L G Q Q

70 80 90 100 110 120
gtgggttcgcatttcctgttgccctcctgccggggagcggccgctgctgggcgagcgc
V G S H F L L P P A G E R P P L L G E R

130 140 150 160 170 180
aggagcgcggcggagcggagcgcgcgcggcggggcggggctgcgagctggcgacctg
R S A A E R S A R G G P G A A Q L A H L

190 200 210 220 230 240
cacggcatcctgcgcccgccgagctctattgccgcaccggcttcacctgcagatcctg
H G I L R R R Q L Y C R T G F H L Q I L

250 260 270 280 290 300
cccgcggcagcagtgccagggcacccggcaggaccacagcctcttcggtatcttgggaattc
P D G S V Q G T R Q D H S L F G I L E F

310 320 330 340 350 360
atcagtgtggcagtgaggactggtcagtattagaggtgtggacagtggtctctatcttggg
I S V A V G L V S I R G V D S G L Y L G

370 380 390 400 410 420
atgaatgacaaaggagaactctatggatcagagaaacttactccgaatgcattctttagg
M N D K G E L Y G S E K L T S E C I F R

430 440 450 460 470 480
gagcagtttgaagagaactgggtataacacctattcatctaacaatataaacatggagac
E Q F E E N W Y N T Y S S N I Y K H G D

490 500 510 520 530 540
actggccgcaggtattttgtggcacttaacaaagacggaactccaagagatggcgccagg
T G R R Y F V A L N K D G T P R D G A R

550 560 570 580 590 600
tccaagaggcatcagaaatttacacatttcttacctagaccagtggtatccagaaagagtt
S K R H Q K F T H F L P R P V D P E R V

610 620 630 640
ccagaattgtacaaggacctactgatgtacacttga
P E L Y K D L L M Y T *

000201" 54626960

[illegible]

AmAcid	Codon	Number	/1000	Fraction
Gly	GGG	13.00	1.89	0.02
Gly	GGA	3.00	0.44	0.00
Gly	GGU	365.00	52.99	0.59
Gly	GGC	238.00	34.55	0.38
Glu	GAG	108.00	15.68	0.22
Glu	GAA	394.00	57.20	0.78
Asp	GAU	149.00	21.63	0.33
Asp	GAC	298.00	43.26	0.67
Val	GUG	93.00	13.50	0.16
Val	GUA	146.00	21.20	0.26
Val	GUU	289.00	41.96	0.51
Val	GUC	38.00	5.52	0.07
Ala	GCG	161.00	23.37	0.26
Ala	GCA	173.00	25.12	0.28
Ala	GCU	212.00	30.78	0.35
Ala	GCC	62.00	9.00	0.10
Arg	AGG	1.00	0.15	0.00
Arg	AGA	0.00	0.00	0.00
Ser	AGU	9.00	1.31	0.03
Ser	AGC	71.00	10.31	0.20
Lys	AAG	111.00	16.11	0.26
Lys	AAA	320.00	46.46	0.74
Asn	AAU	19.00	2.76	0.06
Asn	AAC	274.00	39.78	0.94
Met	AUG	170.00	24.68	1.00
Ile	AUA	1.00	0.15	0.00
Ile	AUU	70.00	10.16	0.17
Ile	AUC	345.00	50.09	0.83
Thr	ACG	25.00	3.63	0.07
Thr	ACA	14.00	2.03	0.04

Figure 10 (continued)

AmAcid	Codon	Number	/1000	Fraction
Thr	ACU	130.00	18.87	0.35
Thr	ACC	206.00	29.91	0.55
Trp	UGG	55.00	7.98	1.00
End	UGA	0.00	0.00	0.00
Cys	UGU	22.00	3.19	0.49
Cys	UGC	23.00	3.34	0.51
End	UAG	0.00	0.00	0.00
End	UAA	0.00	0.00	0.00
Tyr	UAU	51.00	7.40	0.25
Tyr	UAC	157.00	22.79	0.75
Leu	UUG	18.00	2.61	0.03
Leu	UUA	12.00	1.74	0.02
Phe	UUU	51.00	7.40	0.24
Phe	UUC	166.00	24.10	0.76
Ser	UCG	14.00	2.03	0.04
Ser	UCA	7.00	1.02	0.02
Ser	UCU	120.00	17.42	0.34
Ser	UCC	131.00	19.02	0.37
Arg	CGG	1.00	0.15	0.00
Arg	CGA	2.00	0.29	0.01
Arg	CGU	290.00	42.10	0.74
Arg	CGC	96.00	13.94	0.25
Gln	CAG	233.00	33.83	0.86
Gln	CAA	37.00	5.37	0.14
His	CAU	18.00	2.61	0.17
His	CAC	85.00	12.34	0.83
Leu	CUG	480.00	69.69	0.83
Leu	CUA	2.00	0.29	0.00
Leu	CUU	25.00	3.63	0.04
Leu	CUC	38.00	5.52	0.07
Pro	CCG	190.00	27.58	0.77
Pro	CCA	36.00	5.23	0.15
Pro	CCU	19.00	2.76	0.08
Pro	CCC	1.00	0.15	0.00

Figure 11

Codon usage for yeast (highly expressed) genes

AmAcid	Codon	Number	/1000	Fraction	..
Gly	GGG	33.00	0.86	0.01	
Gly	GGA	70.00	1.82	0.02	
Gly	GGT	2672.00	69.62	0.91	
Gly	GGC	171.00	4.46	0.06	
Glu	GAG	277.00	7.22	0.10	
Glu	GAA	2442.00	63.63	0.90	
Asp	GAT	1100.00	28.66	0.48	
Asp	GAC	1211.00	31.55	0.52	
Val	GTG	117.00	3.05	0.04	
Val	GTA	75.00	1.95	0.03	
Val	GTT	1548.00	40.33	0.56	
Val	GTC	1026.00	26.73	0.37	
Ala	GCG	36.00	0.94	0.01	
Ala	GCA	203.00	5.29	0.06	
Ala	GCT	2221.00	57.87	0.65	
Ala	GCC	969.00	25.25	0.28	
Arg	AGG	20.00	0.52	0.01	
Arg	AGA	1336.00	34.81	0.83	
Ser	AGT	116.00	3.02	0.05	
Ser	AGC	94.00	2.45	0.04	
Lys	AAG	2365.00	61.62	0.78	
Lys	AAA	651.00	16.96	0.22	
Asn	AAT	347.00	9.04	0.22	
Asn	AAC	1259.00	32.80	0.78	
Met	ATG	766.00	19.96	1.00	
Ile	ATA	43.00	1.12	0.02	
Ile	ATT	1223.00	31.87	0.52	
Ile	ATC	1070.00	27.88	0.46	

0969945-102000

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

Thr	ACG	28.00	0.73	0.01
Thr	ACA	126.00	3.28	0.06
Thr	ACT	1129.00	29.42	0.50
Thr	ACC	962.00	25.07	0.43
Trp	TGG	325.00	8.47	1.00
End	TGA	10.00	0.26	0.09
Cys	TGT	254.00	6.62	0.89
Cys	TGC	33.00	0.86	0.11
End	TAG	11.00	0.29	0.10
End	TAA	85.00	2.21	0.80
Tyr	TAT	219.00	5.71	0.19
Tyr	TAC	913.00	23.79	0.81
Leu	TTG	2202.00	57.38	0.69
Leu	TTA	576.00	15.01	0.18
Phe	TTT	432.00	11.26	0.27
Phe	TTC	1145.00	29.83	0.73
Ser	TCG	26.00	0.68	0.01
Ser	TCA	149.00	3.88	0.06
Ser	TCT	1279.00	33.33	0.52
Ser	TCC	818.00	21.31	0.33
Arg	CGG	0.00	0.00	0.00
Arg	CGA	1.00	0.03	0.00
Arg	CGT	249.00	6.49	0.15
Arg	CGC	5.00	0.13	0.00
Gln	CAG	62.00	1.62	0.05
Gln	CAA	1225.00	31.92	0.95
His	CAT	236.00	6.15	0.35
His	CAC	433.00	11.28	0.65
Leu	CTG	52.00	1.35	0.02
Leu	CTA	236.00	6.15	0.07
Leu	CTT	90.00	2.35	0.03
Leu	CTC	14.00	0.36	0.00

Figure 12

Codon usage for *Drosophila* (highly expressed) genes

AmAcid	Codon	Number	/1000	Fraction	..
Gly	GGG	6.00	0.28	0.00	
Gly	GGA	380.00	18.04	0.22	
Gly	GGT	575.00	27.29	0.34	
Gly	GGC	746.00	35.41	0.44	
Glu	GAG	1217.00	57.77	0.91	
Glu	GAA	115.00	5.46	0.09	
Asp	GAT	503.00	23.88	0.43	
Asp	GAC	654.00	31.04	0.57	
Val	GTG	719.00	34.13	0.45	
Val	GTA	29.00	1.38	0.02	
Val	GTT	226.00	10.73	0.14	
Val	GTC	608.00	28.86	0.38	
Ala	GCG	94.00	4.46	0.05	
Ala	GCA	80.00	3.80	0.04	
Ala	GCT	446.00	21.17	0.24	
Ala	GCC	1277.00	60.61	0.67	
Arg	AGG	48.00	2.28	0.06	
Arg	AGA	12.00	0.57	0.01	
Ser	AGT	16.00	0.76	0.01	
Ser	AGC	267.00	12.67	0.23	
Lys	AAG	1360.00	64.55	0.93	
Lys	AAA	108.00	5.13	0.07	
Asn	AAT	127.00	6.03	0.13	
Asn	AAC	878.00	41.67	0.87	
Met	ATG	387.00	18.37	1.00	
Ile	ATA	4.00	0.19	0.00	
Ile	ATT	390.00	18.51	0.29	
Ile	ATC	969.00	45.99	0.71	

Figure 12 (continued)

Thr	ACG	114.00	5.41	0.08
Thr	ACA	34.00	1.61	0.02
Thr	ACT	164.00	7.78	0.11
Thr	ACC	1127.00	53.49	0.78
Trp	TGG	243.00	11.53	1.00
End	TGA	1.00	0.05	0.01
Cys	TGT	20.00	0.95	0.08
Cys	TGC	220.00	10.44	0.92
End	TAG	12.00	0.57	0.17
End	TAA	58.00	2.75	0.82
Tyr	TAT	113.00	5.36	0.16
Tyr	TAC	574.00	27.25	0.84
Leu	TTG	210.00	9.97	0.12
Leu	TTA	9.00	0.43	0.01
Phe	TTT	62.00	2.94	0.09
Phe	TTC	635.00	30.14	0.91
Ser	TCG	195.00	9.26	0.17
Ser	TCA	29.00	1.38	0.02
Ser	TCT	103.00	4.89	0.09
Ser	TCC	558.00	26.49	0.48
Arg	CGG	7.00	0.33	0.01
Arg	CGA	25.00	1.19	0.03
Arg	CGT	281.00	13.34	0.34
Arg	CGC	465.00	22.07	0.55
Gln	CAG	703.00	33.37	0.91
Gln	CAA	66.00	3.13	0.09
His	CAT	88.00	4.18	0.22
His	CAC	312.00	14.81	0.78
Leu	CTG	1182.00	56.10	0.69
Leu	CTA	21.00	1.00	0.01
Leu	CTT	55.00	2.61	0.03
Leu	CTC	224.00	10.63	0.13

00000T"54626960

[illegible]

Pro	CCG	84.00	3.99	0.09
Pro	CCA	135.00	6.41	0.15
Pro	CCT	72.00	3.42	0.08
Pro	CCC	626.00	29.71	0.68

DECLARATION AND POWER OF ATTORNEY

As the below-named inventors, we declare that:

Our residences, post office addresses, and citizenships are as stated below under our names.

We believe we are the original, first, and joint inventors of the invention entitled "HUMAN FGF-20 GENE AND GENE EXPRESSION PRODUCTS," which is described and claimed in the foregoing specification and for which a patent is sought.

We hereby claim the benefit under Title 35, United States Code, §120 and/or 119(e) of any United States application(s) and/or provisional application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

Application Serial Number	Filing Date	Status Patented, Pending, Abandoned
60/161,162	10/22/99	Pending
60/187,856	03/08/00	Pending

We have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment specifically referred to herein (if any).

We acknowledge our duty to disclose information of which we are aware which is material to the patentability and examination of this application in accordance with 37 C.F.R. § 1.56(a).

We hereby appoint RICHARD W. SEED, Reg. No. 16,557; ROBERT J. BAYNHAM, Reg. No. 22,846; GEORGE C. RONDEAU, JR., Reg. No. 28,893; DAVID H. DEITS, Reg. No. 28,066; WILLIAM O. FERRON, JR., Reg. No. 30,633; DAVID J. MAKI, Reg. No. 31,392; RICHARD G. SHARKEY, Reg. No. 32,629; DAVID V. CARLSON, Reg. No. 31,153; KARL R. HERMANN, Reg. No. 33,507; DAVID D. MCMASTERS, Reg. No. 33,963; MICHAEL J. DONOHUE, Reg. No. 35,859; JANE E. R. POTTER, Reg. No. 33,332; ROBERT IANNUCCI, Reg. No. 33,514; LORRAINE LINFORD, Reg. No. 35,939; DAVID W. PARKER, Reg. No. 37,414; E. RUSSELL TARLETON, Reg. No. 31,800; Ellen M. Bierman, Reg. No. 38,079; BRIAN G. BODINE, Reg. No. 40,520; BRYAN C. OGDEN, REG. No. 25,362; ANN T. KADLECEK, Reg. No. 39,244; KEVIN S. COSTANZA, Reg. No. 37,801;

THOMAS E. LOOP, Reg. No. 42,810; STEPHEN J. ROSENMAN, Reg. No. 43,058; Brian L. Johnson, Reg. No. 40,033; SUSAN D. BETCHER, Reg. No. 43,498; WILLIAM T. CHRISTIANSEN, Reg. No. 44,614, GARY M. MYLES, Reg. No. 46,209; ERIC J. GASH, Reg. No. 46,274; and JEFFREY C. PEPE, Reg. No. P-46,985; comprising the firm of Seed Intellectual Property Law Group PLLC, 701 Fifth Avenue, Suite 6300, Seattle, Washington 98104-7092, as our attorneys to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. **Please direct all telephone calls to Jane E. R. Potter at (206) 622-4900 and telecopies to (206) 682-6031.**

We further declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Nobuyuki Itoh

Date _____

Residence : City of Kyoto,
Country of Japan
Citizenship : Japan
P.O. Address : Yoshida-Shimoadachi
Sakyo
Kyoto, Japan 606-8501

W. Michael Kavanaugh

Date _____

Residence : City of Mill Valley,
State of California
Citizenship : United States
P.O. Address : c/o Chiron Corporation
P.O. Box 8097
Emeryville, CA 94662-8097

U: CarolW\Chiron\201130\407\Application filing